

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 February 2003 (13.02.2003)

PCT

(10) International Publication Number
WO 03/011838 A1

(51) International Patent Classification⁷: **C07D 239/28**,
239/48, A61K 31/506, A61P 35/00, 19/02

(74) Common Representative: **MERCK & CO., INC.**; 126
East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(21) International Application Number: **PCT/US02/23882**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(22) International Filing Date: **27 July 2002 (27.07.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/309,407 **1 August 2001 (01.08.2001)** **US**

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRALEY, Mark, E.**
[US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-
0907 (US). **HOFFMAN, William, F.** [US/US]; 126 East
Lincoln Avenue, Rahway, NJ 07065-0907 (US). **HART-**
MAN, George, D. [US/US]; 126 East Lincoln Avenue,
Rahway, NJ 07065-0907 (US).

Published:

— *with international search report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 03/011838 A1

(54) Title: **TYROSINE KINASE INHIBITORS**

(57) Abstract: The present invention relates to compounds which inhibit, regulate and/or modulate tyrosine kinase signal transduction, compositions which contain these compounds, and methods of using them to treat tyrosine kinase-dependent diseases and conditions, such as angiogenesis, cancer, tumor growth, atherosclerosis, age related macular degeneration, diabetic retinopathy, inflammatory diseases, and the like in mammals.

TITLE OF THE INVENTION
TYROSINE KINASE INHIBITORS

BACKGROUND OF THE INVENTION

5 The present invention relates to compounds which inhibit, regulate and/or modulate tyrosine kinase signal transduction, compositions which contain these compounds, and methods of using them to treat tyrosine kinase-dependent diseases and conditions, such as angiogenesis, cancer, tumor growth, atherosclerosis, age related macular degeneration, diabetic retinopathy, inflammatory diseases, and
10 the like in mammals.

 Tyrosine kinases are a class of enzymes that catalyze the transfer of the terminal phosphate of adenosine triphosphate to tyrosine residues in protein substrates. Tyrosine kinases play critical roles in signal transduction for a number of cell functions. Though the exact mechanisms of signal transduction is still
15 unclear, tyrosine kinases have been shown to be important contributing factors in cell proliferation, carcinogenesis and cell differentiation.

 Tyrosine kinases can be categorized as receptor type or non-receptor type. Receptor type tyrosine kinases have an extracellular, a transmembrane, and an intracellular portion, while non-receptor type tyrosine kinases are wholly intracellular.

20 The receptor-type tyrosine kinases are comprised of a large number of transmembrane receptors with diverse biological activity. In fact, about twenty different subfamilies of receptor-type tyrosine kinases have been identified. One tyrosine kinase subfamily, designated the HER subfamily, is comprised of EGFR, HER2, HER3, and HER4. Ligands of this subfamily of receptors include epithelial growth factor, TGF- α , amphiregulin, HB-EGF, betacellulin and heregulin. Another
25 subfamily of these receptor-type tyrosine kinases is the insulin subfamily, which includes INS-R, IGF-IR, and IR-R. The PDGF subfamily includes the PDGF- α and β receptors, CSFIR, c-kit and FLK-II. Then there is the FLK family which is comprised of the kinase insert domain receptor (KDR), fetal liver kinase-1 (FLK-1), fetal liver
30 kinase-4 (FLK-4) and the fms-like tyrosine kinase-1 (flt-1). The PDGF and FLK families are usually considered together due to the similarities of the two groups. For a detailed discussion of the receptor-type tyrosine kinases, see Plowman et al., *DN&P* 7(6):334-339, 1994, which is hereby incorporated by reference.

The non-receptor type of tyrosine kinases is also comprised of numerous subfamilies, including Src, Frk, Btk, Csk, Abl, Zap70, Fes/Fps, Fak, Jak, Ack, and LIMK. Each of these subfamilies is further sub-divided into varying receptors. For example, the Src subfamily is one of the largest and includes Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr, and Yrk. The Src subfamily of enzymes has been linked to oncogenesis. For a more detailed discussion of the non-receptor type of tyrosine kinases, see Bolen *Oncogene*, 8:2025-2031 (1993), which is hereby incorporated by reference.

Both receptor-type and non-receptor type tyrosine kinases are implicated in cellular signaling pathways leading to numerous pathogenic conditions, including cancer, psoriasis and hyperimmune responses.

Several receptor-type tyrosine kinases, and the growth factors that bind thereto, have been suggested to play a role in angiogenesis, although some may promote angiogenesis indirectly (Mustonen and Alitalo, *J. Cell Biol.* 129:895-898, 1995). One such receptor-type tyrosine kinase is fetal liver kinase 1 or FLK-1. The human analog of FLK-1 is the kinase insert domain-containing receptor KDR, which is also known as vascular endothelial cell growth factor receptor 2 or VEGFR-2, since it binds VEGF with high affinity. Finally, the murine version of this receptor has also been called NYK (Oelrichs et al., *Oncogene* 8(1):11-15, 1993). VEGF and KDR are a ligand-receptor pair that play an important role in the proliferation of vascular endothelial cells, and the formation and sprouting of blood vessels, termed vasculogenesis and angiogenesis, respectively.

Angiogenesis is characterized by excessive activity of vascular endothelial growth factor (VEGF). VEGF is actually comprised of a family of ligands (Klagsburn and D'Amore, *Cytokine & Growth Factor Reviews* 7:259-270, 1996). VEGF binds the high affinity membrane-spanning tyrosine kinase receptor KDR and the related fms-like tyrosine kinase-1, also known as Flt-1 or vascular endothelial cell growth factor receptor 1 (VEGFR-1). Cell culture and gene knockout experiments indicate that each receptor contributes to different aspects of angiogenesis. KDR mediates the mitogenic function of VEGF whereas Flt-1 appears to modulate non-mitogenic functions such as those associated with cellular adhesion. Inhibiting KDR thus modulates the level of mitogenic VEGF activity. In fact, tumor growth has been shown to be susceptible to the antiangiogenic effects of VEGF receptor antagonists. (Kim et al., *Nature* 362, pp. 841-844, 1993).

Solid tumors can therefore be treated by tyrosine kinase inhibitors since these tumors depend on angiogenesis for the formation of the blood vessels necessary to support their growth. These solid tumors include histiocytic lymphoma, cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung, including lung adenocarcinoma and small cell lung cancer. Additional examples include cancers in which overexpression or activation of Raf-activating oncogenes (e.g., K-ras, erb-B) is observed. Such cancers include pancreatic and breast carcinoma. Accordingly, inhibitors of these tyrosine kinases are useful for the prevention and treatment of proliferative diseases dependent on these enzymes.

10 The angiogenic activity of VEGF is not limited to tumors. VEGF accounts for most of the angiogenic activity produced in or near the retina in diabetic retinopathy. This vascular growth in the retina leads to visual degeneration culminating in blindness. Ocular VEGF mRNA and protein are elevated by conditions such as retinal vein occlusion in primates and decreased pO₂ levels in mice that lead to
15 neovascularization. Intraocular injections of anti-VEGF monoclonal antibodies or VEGF receptor immunofusions inhibit ocular neovascularization in both primate and rodent models. Regardless of the cause of induction of VEGF in human diabetic retinopathy, inhibition of ocular VEGF is useful in treating the disease.

 Expression of VEGF is also significantly increased in hypoxic regions
20 of animal and human tumors adjacent to areas of necrosis. VEGF is also upregulated by the expression of the oncogenes ras, raf, src and mutant p53 (all of which are relevant to targeting cancer). Monoclonal anti-VEGF antibodies inhibit the growth of human tumors in nude mice. Although these same tumor cells continue to express VEGF in culture, the antibodies do not diminish their mitotic rate. Thus tumor-
25 derived VEGF does not function as an autocrine mitogenic factor. Therefore, VEGF contributes to tumor growth in vivo by promoting angiogenesis through its paracrine vascular endothelial cell chemotactic and mitogenic activities. These monoclonal antibodies also inhibit the growth of typically less well vascularized human colon cancers in athymic mice and decrease the number of tumors arising from inoculated
30 cells.


 Viral expression of a VEGF-binding construct of Flk-1, Flt-1, the mouse KDR receptor homologue, truncated to eliminate the cytoplasmic tyrosine kinase domains but retaining a membrane anchor, virtually abolishes the growth of a transplantable glioblastoma in mice presumably by the dominant negative mechanism
35 of heterodimer formation with membrane spanning endothelial cell VEGF receptors.

Embryonic stem cells, which normally grow as solid tumors in nude mice, do not produce detectable tumors if both VEGF alleles are knocked out. Taken together, these data indicate the role of VEGF in the growth of solid tumors. Inhibition of KDR or Flt-1 is implicated in pathological angiogenesis, and these receptors are
 5 useful in the treatment of diseases in which angiogenesis is part of the overall pathology, e.g., inflammation, diabetic retinal vascularization, as well as various forms of cancer since tumor growth is known to be dependent on angiogenesis. (Weidner et al., N. Engl. J. Med., 324, pp. 1-8, 1991).

Accordingly, the identification of small compounds which specifically
 10 inhibit, regulate and/or modulate the signal transduction of tyrosine kinases is desirable and is an object of this invention.

SUMMARY OF THE INVENTION


The present invention relates to compounds that are capable of
 15 inhibiting, modulating and/or regulating signal transduction of both receptor-type and non-receptor type tyrosine kinases. One embodiment of the present invention is illustrated by a compound of Formula I, and the pharmaceutically acceptable salts thereof:



20

DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of kinases and are illustrated by a compound of Formula I:



wherein

R^{1a} is independently selected from:

- 5 (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- (3) OR⁸, and
- (4) N(R⁸)₂;

10 R^1 is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- (3) unsubstituted or substituted C₃-C₁₀ cycloalkyl,
- (4) unsubstituted or substituted aryl,
- 15 (5) unsubstituted or substituted heterocycle,
- (6) halo,
- (7) CF₃,
- (8) -(CH₂)_t R⁹C(O)R⁸,
- (9) -C(O)R⁹,
- 20 (10) -(CH₂)_tOR⁸,
- (11) unsubstituted or substituted C₂-C₆ alkenyl,
- (12) unsubstituted or substituted C₂-C₆ alkynyl,
- (13) CN,
- (14) -(CH₂)_t NR⁷ R⁸,
- 25 (15) -(CH₂)_t C(O)NR⁷R⁸,
- (16) -C(O)OR⁸, and
- (17) -(CH₂)_t S(O)_q(CH₂)_tNR⁷R⁸;

R² is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- 5 (3) unsubstituted or substituted C₃-C₁₀ cycloalkyl,
- (4) unsubstituted or substituted aryl,
- (5) unsubstituted or substituted heterocycle,
- (6) halo,
- (7) CF₃,
- 10 (8) -(CH₂)_t R⁹C(O)R⁸,
- (9) -C(O)R⁹,
- (10) -(CH₂)_tOR⁸,
- (11) unsubstituted or substituted C₂-C₆ alkenyl,
- (12) unsubstituted or substituted C₂-C₆ alkynyl,
- 15 (13) CN,
- (14) -(CH₂)_t NR⁷ R⁸,
- (15) -(CH₂)_t C(O)NR⁷R⁸,
- (16) -C(O)OR⁸, and
- (17) -(CH₂)_t S(O)_q(CH₂)_tNR⁷R⁸

R³ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- (3) unsubstituted or substituted aralkyl,
- 25 (4) CN,
- (5) halo,
- (6) N(R⁸)₂,
- (7) OR⁸, and
- (8) unsubstituted or substituted aryl;

R⁷ is selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
- (3) unsubstituted or substituted aralkyl;

R⁸ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- 5 (3) unsubstituted or substituted aryl,
- (4) unsubstituted or substituted heterocycle,
- (5) unsubstituted or substituted C₃-C₁₀ cycloalkyl, and
- (6) unsubstituted or substituted aralkyl;

- 10 R⁷ and R⁸, when attached to the same nitrogen atom may be joined to form a 5-7 membered heterocycle containing, in addition to the nitrogen, one or two more heteroatoms selected from N, O, or S, said heterocycle being optionally substituted with one to three R² substituents;

- 15 R⁹ is independently selected from:

- (1) unsubstituted or substituted C₁-C₁₀ alkyl,
- (2) unsubstituted or substituted heterocycle, and
- (3) unsubstituted or substituted aryl;

- 20 W is selected from:

- (1) aryl, and
- (2) heterocycle;

m is 0, 1, or 2;

- 25 n is independently 0, 1, 2, 3, 4, 5, or 6;


p is 0, 1, 2, 3, or 4;

q is independently 0, 1, or 2;

t is independently 0, 1, 2, 3, 4, 5, or 6;

- 30 or a pharmaceutically acceptable salt, hydrate or stereoisomer thereof.

Another embodiment of the instant invention is illustrated by a compound of Formula I:



wherein

R^{1a} is independently selected from:

- 5 (1) H,
- (2) unsubstituted or substituted C_1 - C_6 alkyl, and
- (3) OR^8 ;

R^1 is independently selected from:

- 10 (1) H,
- (2) unsubstituted or substituted C_1 - C_{10} alkyl,
- (3) halo,
- (4) CF_3 ,
- (5) $-(CH_2)_t R^9 C(O)R^8$,
- 15 (6) $-C(O)R^9$,
- (7) $-(CH_2)_t OR^8$,
- (8) $-(CH_2)_t C(O)NR^7 R^8$,
- (9) $-C(O)OR^8$,
- (10) $-(CH_2)_t NR^7 R^8$, and
- 20 (11) $-(CH_2)_t S(O)_q (CH_2)_t NR^7 R^8$;

R^2 is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C_1 - C_{10} alkyl,
- 25 (3) unsubstituted or substituted aryl,
- (4) unsubstituted or substituted heterocycle,

- 5
- (5) unsubstituted or substituted C₃-C₁₀ cycloalkyl,
 - (6) unsubstituted or substituted C₂-C₆ alkenyl,
 - (7) unsubstituted or substituted C₂-C₆ alkynyl,
 - (8) CN,
 - (9) halo,
 - (10) N(R⁸)₂, and
 - (11) OR⁸;

R³ is independently selected from:

- 10
- (1) H,
 - (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
 - (3) unsubstituted or substituted aralkyl;

R⁷ is selected from:

- 15
- (1) H,
 - (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
 - (3) unsubstituted or substituted aralkyl;

R⁸ is independently selected from:

- 20
- (1) H,
 - (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
 - (3) unsubstituted or substituted aryl;

25 R⁷ and R⁸, when attached to the same nitrogen atom may be joined to form a 5-7 membered heterocycle containing, in addition to the nitrogen, one or two more heteroatoms selected from N, O, or S, said heterocycle being optionally substituted with one to three R² substituents;

R⁹ is independently selected from

- 30
- (1) unsubstituted or substituted aryl, and
 - (2) unsubstituted or substituted heterocycle;

W is selected from:

- (1) aryl, and

(2) heteroaryl, selected from pyridyl, pyrimidinyl, isoxazolyl, or pyrazinyl;

m is 0, 1, or 2;

n is independently 0, 1, 2, 3, 4, 5, or 6;

5 p is 0, 1, 2, 3, or 4;


q is independently 0, 1, or 2;

t is independently 0, 1, 2, 3, 4, 5, or 6;

or a pharmaceutically acceptable salt, hydrate or stereoisomer thereof.

10

Another embodiment of the instant invention is illustrated by a compound of Formula II:



15

wherein

R¹ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- 20 (3) halo,
- (4) unsubstituted or substituted aryl,
- (5) unsubstituted or substituted heterocycle,
- (6) CF₃,
- (7) -(CH₂)_t R⁹C(O)R⁸,
- 25 (8) -C(O)R⁹, and
- (9) -(CH₂)_tOR⁸;

R² is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- (3) unsubstituted or substituted aryl,
- 5 (4) unsubstituted or substituted heterocycle,
- (5) halo,
- (6) OR⁸,
- (7) N(R⁸)₂, and
- (8) CN;

10

R³ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
- 15 (3) unsubstituted or substituted aralkyl;

15

R⁸ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
- 20 (3) unsubstituted or substituted aryl;

20

R⁹ is independently selected from

- (1) unsubstituted or substituted aryl, and
- (2) unsubstituted or substituted heterocycle;

25 m is 0, 1, or 2;

n is 0, 1, 2, 3, 4, 5, or 6;

p is 0, 1, 2, 3, or 4;

t is independently 0, 1, 2, 3, 4, 5, or 6;

30 or a pharmaceutically acceptable salt, hydrate or stereoisomer thereof.

Examples of compounds of the instant invention include

4-(2-amino-5-bromo-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine;


4-(2-amino-1,3-thiazol-4-yl)-N-(3,5dimethylphenyl)pyrimidin-2-amine;

4-(2-amino-5-phenyl-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine;
 2-amino-4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-
 carbonitrile;
 4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile;


5

or a pharmaceutically acceptable salt, or hydrate thereof.

Specific examples of compounds of the instant invention include



10 2-amino-4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-
 carbonitrile;



4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile;

15 or a pharmaceutically acceptable salt, or hydrate thereof.

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-
 20 1190), and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, being

included in the present invention. In addition, the compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be encompassed by the scope of the invention, even though only one tautomeric structure is depicted.


When any variable (e.g. aryl, heterocycle, R¹, R² etc.) occurs more
5 than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds.

Lines drawn into the ring systems from substituents (such as from
R², R³, etc.) indicate that the indicated bond may be attached to any of the
10 substitutable ring carbon atoms or heteroatoms. If the ring system is polycyclic, it is intended that the bond be attached to any of the suitable carbon atoms or heteroatoms on the proximal ring only.

It is understood that substituents and substitution patterns on the
compounds of the instant invention can be selected by one of ordinary skill in the art
15 to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials.

As used herein, "alkyl" is intended to include both branched and
straight-chain hydrocarbon groups having the specified number of carbon atoms. For
20 example, C₁-C₁₀, as in "C₁-C₁₀ alkyl" is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear or branched arrangement. For example, "C₁-C₁₀ alkyl" specifically includes methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, adamantyl, and so on.

"Cycloalkyl" as used herein is intended to include non-aromatic cyclic
25 hydrocarbon groups, having the specified number of carbon atoms, which may or may not be bridged or structurally constrained. Examples of such cycloalkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, cyclooctyl, cycloheptyl, tetrahydro-naphthalene, methylenecyclohexyl, and the like. As used herein, examples of "C₃ - C₁₀ cycloalkyl" may include, but are not limited to:



As used herein, the term "alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge.

- 5 If no number of carbon atoms is specified, the term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to 4 non-aromatic carbon-carbon double bonds may be present. Thus, "C₂-C₆ alkenyl" means an alkenyl radical
- 10 having from 2 to 6 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

- The term "alkynyl" refers to a hydrocarbon radical straight, branched
- 15 or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to 3 carbon-carbon triple bonds may be present. Thus, "C₂-C₆ alkynyl" means an alkynyl radical having from 2 to 6 carbon atoms. Alkynyl groups include ethynyl, propynyl and butynyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkynyl group may contain triple bonds and
- 20 may be substituted if a substituted alkynyl group is indicated.

- As used herein, "aryl" is intended to mean any stable monocyclic, bicyclic or tricyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydro-
- 25 naphthyl, indanyl, indanonyl, biphenyl, tetralinyl, tetralonyl, fluorenonyl, phenanthryl, anthryl, acenaphthyl, dihydroindanyl, dihydroindenyl, and the like.

As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro, fluoro, bromo and iodo.

The term "heteroaryl", as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to: acridinyl, benzodioxolyl, benzofuranyl, benzofurazanyl, benzoimidazolyl, benzopyranyl, benzopyrazolyl, benzotriazolyl, benzothiazolyl, benzothienyl, benzothiofuranyl, benzothiophenyl, benzothiopyranyl, benzoxazolyl, carbazolyl, cinnolinyl, imidazolyl, imidazolyl, imidazolidinyl, indazolyl, quinoxalyl, isobenzofuranyl, pyrazolyl, indolyl, furanyl, thienyl, quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, tetrahydroquinoline, triazolyl, and the like.

The term heterocycle or heterocyclic or heterocyclyl, as used herein, represents a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. "Heterocycle" or "heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrathydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: benzodioxolyl, benzofuranyl, benzofurazanyl, benzoimidazolyl, benzopyranyl, benzopyrazolyl, benzotriazolyl, benzothiazolyl, benzothienyl, benzothiofuranyl, benzothiophenyl, benzothiopyranyl, benzoxazolyl, carbazolyl, carbolinyl, chromanyl, cinnolinyl, dihydrobenzofuranyl, dihydrobenzofuryl, dihydrobenzoimidazolyl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl,

dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl,
 dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolinyl, dihydrotetrazolyl,
 dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl,
 dihydroazetidiny, furyl, furanyl, imidazolyl, imidazoliny, imidazolidinyl, indazolyl,
 5 indolaziny, indolinyl, indolyl, isobenzofuranyl, isochromanyl, isoindolyl,
 isoindolinyl, isoquinolinone, isoquinolyl, isothiazolyl, isothiazolidinyl, isoxazoliny, isoxazolyl,
 methylenedioxybenzoyl, morpholinyl, naphthpyridinyl, oxadiazolyl,
 oxazolyl, oxazoliny, oxetanyl, 2-oxoazepiny, oxadiazolyl, 2-oxopiperazinyl, 2-
 oxopiperdiny, 2-oxopyrrolidinyl, piperidyl, piperidinyl, piperazinyl, pyranyl,
 10 pyrazinyl, pyrazolyl, pyridazinyl, 2-pyridinonyl, pyridopyridinyl, pyridazinyl, pyridyl,
 pyrimidinyl, pyrrolyl, pyrrolidinyl, quinazoliny, quinolinyl, quinolyl, quinolinonyl,
 quinoxaliny, tetrahydrofuryl, tetrahydroisoquinolinyl, tetrahydropyranyl,
 tetrahydroquinolinyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thiazolinyl,
 thienofuryl, thienyl, triazolyl, azetidiny, 1,4-dioxanyl, hexahydroazepiny, and the
 15 like.

As used herein, "aralkyl" is intended to mean an aryl moiety, as
 defined above, attached through a C₁-C₁₀ alkyl linker, where alkyl is defined above.
 Examples of aralkyls include, but are not limited to, benzyl, naphthylmethyl and
 phenylpropyl.


20 As used herein, "heterocyclalkyl" is intended to mean a heteroaryl
 moiety, as defined below, attached through a C₁-C₁₀ alkyl linker, where alkyl is
 defined above. Examples of heterocyclalkyls include, but are not limited to,
 2-pyridylmethyl, 2-imidazolylethyl, 2-quinolinylmethyl, 2-imidazolylmethyl
 and the like.

25 As used herein, the terms "substituted C₁-C₁₀ alkyl" and "substituted
 C₁-C₆ alkoxy" are intended to include the branch or straight-chain alkyl group of the
 specified number of carbon atoms, wherein the carbon atoms may be substituted with
 F, Cl, Br, CF₃, N₃, NO₂, NH₂, oxo, -OH, -O(C₁-C₆ alkyl), S(O)₀₋₂, (C₁-C₆ alkyl)
 S(O)₀₋₂-, (C₁-C₆ alkyl)S(O)₀₋₂(C₁-C₆ alkyl)-, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl,

- C₂-C₆ alkynyl, -C(O)NH, (C₁-C₆ alkyl) C(O)NH-, H₂NC(NH)-, (C₁-C₆ alkyl)C(O)-, -O(C₁-C₆ alkyl)CF₃, (C₁-C₆ alkyl)OC(O)-, (C₁-C₆ alkyl)O(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)C(O)₂(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)OC(O)NH-, aryl, benzyl, heterocycle, aralkyl, heterocyclalkyl, halo-aryl, halo-benzyl, halo-heterocycle, cyano-aryl, cyano-benzyl and cyano-heterocycle.

- As used herein, the terms "substituted aryl", "substituted heterocycle", "substituted aralkyl" and "substituted heterocyclalkyl" are intended to include the cyclic group containing from 1 to 3 substituents in addition to the point of attachment to the rest of the compound. Such substituents are preferably selected from the group which includes but is not limited to F, Cl, Br, CF₃, NH₂, N(C₁-C₆ alkyl)₂, NO₂, CN, N₃, C₁-C₂₀ alkyl, C₁-C₆ alkoxy, -OH, -O(C₁-C₆ alkyl), S(O)₀₋₂, (C₁-C₆ alkyl) S(O)₀₋₂, (C₁-C₆ alkyl)S(O)₀₋₂(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)C(O)NH-, H₂N-C(NH)-, (C₁-C₆ alkyl)C(O)-, (C₁-C₆ alkyl)OC(O)-, (C₁-C₆ alkyl)O(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)C(O)₂(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)OC(O)NH-, aryl, aralkyl, heteroaryl, heterocyclalkyl, halo-aryl, halo-aralkyl, halo-heterocycle, halo-heterocyclalkyl, cyano-aryl, cyano-aralkyl, cyano-heterocycle and cyano-heterocyclalkyl.

The moiety formed when, in the definition of R⁷ and R⁸ are joined to form a ring, is illustrated by, but not limited to, the following:



20

Preferably, R¹ is independently selected from H, unsubstituted or substituted C₁-C₁₀ alkyl, -(CH₂)_tOR⁸, -C(O)R⁹, or halo. Most preferred, R¹ is

independently selected from H, unsubstituted or substituted C₁-C₁₀ alkyl, -(CH₂)_tOR⁸, or halo.

Preferably, R² is independently selected from H, unsubstituted or substituted C₁-C₁₀ alkyl, CN, OR⁸, N(R⁸)₂, halo or unsubstituted or substituted aryl.

5 Most preferred, R² is independently selected from H, unsubstituted or substituted C₁-C₁₀ alkyl, CN, OR⁸, halo, N(R⁸)₂, halo or unsubstituted or substituted aryl.

Preferably, R³ is independently selected from H, or unsubstituted or substituted C₁-C₁₀ alkyl.

Preferably, m is selected from 0 or 1.

10 Preferably, n and p are independently selected from 0, 1, 2 or 3.

Preferably, t is independently selected from 0, 1, 2, 3, or 4.

It is intended that the definition of any substituent or variable (e.g., R², R³, n, etc.) at a particular location in a molecule be independent of its definitions elsewhere in that molecule. Thus, -N(R⁸)₂ represents -NHH, -NHCH₃, -NHC₂H₅,
15 etc. It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials.

20 The pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like: and the
25 salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

The pharmaceutically acceptable salts of the compounds of this invention can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with the appropriate inorganic or organic base.

10

UTILITY

“Tyrosine kinase-dependent diseases or conditions” refers to pathologic conditions that depend on the activity of one or more tyrosine kinases. Tyrosine kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities including proliferation, adhesion and migration, and differentiation. Diseases associated with tyrosine kinase activities include the proliferation of tumor cells, the pathologic neovascularization that supports solid tumor growth, ocular neovascularization (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

20

Included within the scope of the present invention is a pharmaceutical composition which is comprised of a compound of Formula I as described above and a pharmaceutically acceptable carrier. The present invention also encompasses a method of treating or preventing cancer in a mammal in need of such treatment which is comprised of administering to said mammal a therapeutically effective amount of a compound of Formula I. Preferred cancers for treatment are selected from cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung. Another set of preferred forms of cancer are histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.

25

Also included is a method of treating or preventing a disease in which angiogenesis is implicated, which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Formula I. Such a disease in which angiogenesis is implicated is ocular diseases such as retinal vascularization, diabetic retinopathy, age-related macular degeneration, and the like.

Also included is a method of inhibiting at least two tyrosine kinase receptors, selected from KDR, EGFR or SRC, by administering a therapeutically effective amount of a compound of Formula I.

Also included within the scope of the present invention is a method of treating or preventing inflammatory diseases which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Formula I. Examples of such inflammatory diseases are rheumatoid arthritis, psoriasis, contact dermatitis, delayed hypersensitivity reactions, and the like.

Also included is a method of treating or preventing a tyrosine kinase-dependent disease or condition in a mammal which comprises administering to a mammalian patient in need of such treatment a therapeutically effective amount of a compound of Formula I. The therapeutic amount varies according to the specific disease and is discernable to the skilled artisan without undue experimentation.

A method of treating or preventing retinal vascularization which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of compound of Formula I is also encompassed by the present invention. Methods of treating or preventing ocular diseases, such as diabetic retinopathy and age-related macular degeneration, are also part of the invention. Also included within the scope of the present invention is a method of treating or preventing inflammatory diseases, such as rheumatoid arthritis, psoriasis, contact dermatitis and delayed hypersensitivity reactions, as well as treatment or prevention of bone associated pathologies selected from osteosarcoma, osteoarthritis, and rickets.

The invention also contemplates the use of the instantly claimed compounds in combination with a second compound selected from the group consisting of:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) retinoid receptor modulator,
- 4) a cytotoxic agent,
- 5 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor, and
- 10 10) another angiogenesis inhibitor.

The preferred second angiogenesis inhibitor is selected from the group consisting of a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP inhibitor, an integrin blocker, interferon- α , interleukin-12, 15 pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, and an antibody to VEGF. Preferred estrogen receptor modulators are tamoxifen and raloxifene.

Also included in the scope of the claims is a method of treating cancer 20 which comprises administering a therapeutically effective amount of a compound of Formula I in combination with radiation therapy and/or in combination with a compound selected from the group consisting of:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 25 3) retinoid receptor modulator,
- 4) a cytotoxic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 30 8) an HIV protease inhibitor,

9) a reverse transcriptase inhibitor, and

10) another angiogenesis inhibitor.

And yet another embodiment of the invention is a method of treating cancer which comprises administering a therapeutically effective amount of a
5 compound of Formula I in combination with paclitaxel or trastuzumab.

These and other aspects of the invention will be apparent from the teachings contained herein.

The instant compounds are useful as pharmaceutical agents for mammals, especially for humans, in the treatment of tyrosine kinase dependent
10 diseases. Such diseases include the proliferation of tumor cells, the pathologic neovascularization (or angiogenesis) that supports solid tumor growth, ocular neovascularization (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

The compounds of the instant invention may be administered to
15 patients for use in the treatment of cancer. The instant compounds inhibit tumor angiogenesis, thereby affecting the growth of tumors (J. Rak et al. *Cancer Research*, 55:4575-4580, 1995). The anti-angiogenesis properties of the instant compounds are also useful in the treatment of certain forms of blindness related to retinal vascularization.

20 It has been shown that simultaneously targeting multiple angiogenic factors may improve survival of mammals with cancer metastases. (R.M. Shaheen et al., *Cancer Research*, 61:1464-1468, 2001). The disclosed compounds are also useful in a method of treating cancer by administering a therapeutically effective amount of a compound of Formula I in order to inhibit at least two tyrosine kinase
25 receptors, particularly KDR, EGFR and SRC.

The disclosed compounds are also useful in the treatment of certain bone-related pathologies, such as osteosarcoma, osteoarthritis, and rickets, also known as oncogenic osteomalacia. (Hasegawa et al., *Skeletal Radiol.*, 28, pp.41-45, 1999; Gerber et al., *Nature Medicine*, Vol. 5, No. 6, pp.623-628, June 1999). And
30 since VEGF directly promotes osteoclastic bone resorption through KDR/Flk-1

expressed in mature osteoclasts (FEBS Let. 473:161-164 (2000); Endocrinology, 141:1667 (2000)), the instant compounds are also useful to treat and prevent conditions related to bone resorption, such as osteoporosis and Paget's disease.

The claimed compounds can also be used to reduce or prevent tissue damage which occurs after cerebral ischemic events, such as stroke, by reducing cerebral edema, tissue damage, and reperfusion injury following ischemia. (*Drug News Perspect* 11:265-270 (1998); *J. Clin. Invest.* 104:1613-1620 (1999); *Nature Med* 7:222-227 (2001)).

The instant compounds are useful in the treatment of preeclampsia. Studies have shown that the action of VEGF on the Flt-1 receptor is pivotal in the pathogenesis of preeclampsia. (*Laboratory Investigation* 79:1101-1111 (September 1999)). Vessels of pregnant women incubated with VEGF exhibit a reduction in endothelium-dependent relaxation similar to that induced by plasma from women with preeclampsia. In the presence of an anti-Flt-1 receptor antibody, however, neither VEGF or plasma from women with preeclampsia reduced the endothelium-dependent relaxation. Therefore the claimed compounds serve to treat preeclampsia via their action on the tyrosine kinase domain of the Flt-1 receptor.

The instant compounds can also be used to prevent or treat tissue damage during bacterial meningitis, such as tuberculous meningitis. (Matsuyama et al., *J. Neurol. Sci.* 186:75-79 (2001)). The instant invention therefore encompasses a method of treating or preventing tissue damage due to bacterial meningitis which comprises administering a therapeutically effective amount of a compound of Formula 1. Studies have shown that VEGF is secreted by inflammatory cells during bacterial meningitis and that VEGF contributes to blood-brain barrier disruption. (van der Flier et al., *J. Infectious Diseases*, 183:149-153 (2001)). The claimed compounds can inhibit VEGF-induced vascular permeability and therefore serve to prevent or treat blood-brain barrier disruption associated with bacterial meningitis.

The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents, optionally with known adjuvants, such as alum, in a

pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

5 For oral use of a chemotherapeutic compound according to this invention, the selected compound may be administered, for example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and lubricating agents, such as magnesium stearate, are commonly added. For
10 oral administration in capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring agents may be added. For intramuscular, intraperitoneal, subcutaneous and intravenous use, sterile solutions of the active ingredient are usually
15 prepared, and the pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

 The compounds of the instant invention may also be co-administered with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. For example, in the case of bone-related disorders, combinations that would be useful include those with antiresorptive
20 bisphosphonates, such as alendronate and risedronate; integrin blockers (defined further below), such as $\alpha_v\beta_3$ antagonists; conjugated estrogens used in hormone replacement therapy, such as PREMPRO®, PREMARIN® and ENDOMETRION®;
25 selective estrogen receptor modulators (SERMs), such as raloxifene, droloxifene, CP-336,156 (Pfizer) and lasofoxifene; cathepsin K inhibitors; and ATP proton pump inhibitors.

 The instant compounds are also useful in combination with known anti-cancer agents. Such known anti-cancer agents include the following: estrogen
30 receptor modulators, androgen receptor modulators, retinoid receptor modulators,

cytotoxic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, and other angiogenesis inhibitors. The instant compounds are particularly useful when coadministered with radiation therapy. The synergistic effects of inhibiting VEGF in combination with radiation therapy have been described in the art (see WO 00/61186).

“Estrogen receptor modulators” refers to compounds which interfere or inhibit the binding of estrogen to the receptor, regardless of mechanism. Examples of estrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-2H-1-benzopyran-3-yl)-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenyl-hydrazone, and SH646.

“Androgen receptor modulators” refers to compounds which interfere or inhibit the binding of androgens to the receptor, regardless of mechanism. Examples of androgen receptor modulators include finasteride and other 5 α -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole, and abiraterone acetate.

“Retinoid receptor modulators” refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid, α -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl) retinamide, and N-4-carboxyphenyl retinamide.

“Cytotoxic agents” refer to compounds which cause cell death primarily by interfering directly with the cell's functioning or inhibit or interfere with cell myosis, including alkylating agents, tumor necrosis factors, intercalators, microtubulin inhibitors, and topoisomerase inhibitors.

Examples of cytotoxic agents include, but are not limited to, tirapazimine, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate,

trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, profiromycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methylpyridine) platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum (II)]tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplaston, 3'-deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755, and 4-demethoxy-3-deamino-3-aziridinyl-4-methylsulphonyl-daunorubicin (see WO 00/50032).

Examples of microtubulin inhibitors include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vinflunine, cryptophycin, 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl)benzene sulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258, and BMS188797.

Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exo-benzylidene-chartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrazolo[3,4,5-kl]acridine-2-(6H)propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H, 12H-benzo[de]pyrano[3',4':b,7]indolizino[1,2b]quinoline-10,13(9H,15H)dione, lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPI1100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrine, (5a, 5aB, 8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydrofuro(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoguinoline-5,10-dione,


5- (3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrazolo[4,5,1-de]acridin-6-one, N-[1-[2(diethylamino)ethylamino]-7-methoxy-9-oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c]quinolin-7-one, and dimesna.

“Antiproliferative agents” includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emittefur, tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydro-benzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-[2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3H-pyrimidino[5,4-b][1,4]thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-glutamic acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabino furanosyl cytosine, and 3-aminopyridine-2-carboxaldehyde thiosemicarbazone. “Antiproliferative agents” also includes monoclonal antibodies to growth factors, other than those listed under “angiogenesis inhibitors”, such as trastuzumab, and tumor suppressor genes, such as p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S. Patent No. 6,069,134, for example).

“HMG-CoA reductase inhibitors” refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the assays described or cited in U.S. Patent No. 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33. The terms “HMG-CoA reductase inhibitor” and “inhibitor of HMG-CoA reductase” have the same meaning when used herein.

Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR[®]; see U.S. Patent Nos. 4,231,938; 4,294,926; 4,319,039), simvastatin (ZOCOR[®]; see U.S. Patent Nos. 4,444,784; 4,820,850; 4,916,239), pravastatin (PRAVACHOL[®]; see U.S. Patent Nos. 4,346,227; 5 4,537,859; 4,410,629; 5,030,447 and 5,180,589), fluvastatin (LESCOL[®]; see U.S. Patent Nos. 5,354,772; 4,911,165; 4,929,437; 5,189,164; 5,118,853; 5,290,946; 5,356,896), atorvastatin (LIPITOR[®]; see U.S. Patent Nos. 5,273,995; 4,681,893; 5,489,691; 5,342,952) and cerivastatin (also known as rivastatin and BAYCHOL[®]; see US Patent No. 5,177,080). The structural formulae of these and additional HMG-
10 CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the
15 free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase inhibitory activity, and therefor the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention. An illustration of the lactone portion and its corresponding open-acid form is shown below as structures I and II.

20



In HMG-CoA reductase inhibitors where an open-acid form can exist, salt and ester forms may preferably be formed from the open-acid, and all such forms

are included within the meaning of the term "HMG-CoA reductase inhibitor" as used herein. Preferably, the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin, and most preferably simvastatin. Herein, the term "pharmaceutically acceptable salts" with respect to the HMG-CoA reductase inhibitor shall mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base, particularly those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc and tetramethylammonium, as well as those salts formed from amines such as ammonia, ethylenediamine, N-methylglucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, 1-p-chlorobenzyl-2-pyrrolidine-1'-yl-methylbenzimidazole, diethylamine, piperazine, and tris(hydroxymethyl) aminomethane. Further examples of salt forms of HMG-CoA reductase inhibitors may include, but are not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynapthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamaote, palmitate, panthothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide, and valerate.

Ester derivatives of the described HMG-CoA reductase inhibitor compounds may act as prodrugs which, when absorbed into the bloodstream of a warm-blooded animal, may cleave in such a manner as to release the drug form and permit the drug to afford improved therapeutic efficacy.

"Prenyl-protein transferase inhibitor" refers to a compound which inhibits any one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase (FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II,

also called Rab GGPTase). Examples of prenyl-protein transferase inhibiting compounds include (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1*H*)-quinolinone, (-)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1*H*)-quinolinone, (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1*H*)-quinolinone, 5(*S*)-*n*-butyl-1-(2,3-dimethylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, (*S*)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone, 5(*S*)-*n*-Butyl-1-(2-methylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, 1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-2-methyl-5-imidazolylmethyl]-2-piperazinone, 1-(2,2-diphenylethyl)-3-[*N*-(1-(4-cyanobenzyl)-1H-imidazol-5-ylethyl)carbamoyl]piperidine, 4-{5-[4-hydroxymethyl]-4-(4-chloropyridin-2-ylmethyl)-piperidine-1-ylmethyl}-2-methylimidazol-1-ylmethyl} benzonitrile, 4-{5-[4-hydroxymethyl-4-(3-chlorobenzyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl} benzonitrile, 4-{3-[4-(2-oxo-2H-pyridin-1-yl)benzyl]-3H-imidazol-4-ylmethyl} benzonitrile, 4-{3-[4-(5-chloro-2-oxo-2H-[1,2']bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 4-{3-[4-(2-oxo-2H-[1,2']bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 4-[3-(2-oxo-1-phenyl-1,2-dihydropyridin-4-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 18,19-dihydro-19-oxo-5*H*,17*H*-6,10:12,16-dimetheno-1H-imidazo[4,3-*c*][1,11,4]dioxazacyclononadecine-9-carbonitrile, (±)-19,20-dihydro-19-oxo-5*H*-18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxatriaza-cyclooctadecine-9-carbonitrile, 19,20-dihydro-19-oxo-5*H*,17*H*-18,21-ethano-6,10:12,16-dimetheno-22*H*-imidazo[3,4-*h*][1,8,11,14]oxatriazacycloeicosine-9-carbonitrile, and (±)-19,20-dihydro-3-methyl-19-oxo-5*H*-18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxa-triazacyclooctadecine-9-carbonitrile.

Other examples of prenyl-protein transferase inhibitors can be found in the following publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U.S. Patent No. 5,420,245, U.S. Patent No. 5,523,430, U.S. Patent

No. 5,532,359, U.S. Patent No. 5,510,510, U.S. Patent No. 5,589,485, U.S. Patent No. 5,602,098, European Patent Publ. 0 618 221, European Patent Publ. 0 675 112, European Patent Publ. 0 604 181, European Patent Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Patent No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Patent No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U.S. Patent No. 5,532,359. For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see European J. of Cancer, Vol. 35, No. 9, pp.1394-1401 (1999).

Examples of HIV protease inhibitors include amprenavir, abacavir, CGP-73547, CGP-61755, DMP-450, indinavir, nelfinavir, tipranavir, ritonavir, saquinavir, ABT-378, AG 1776, and BMS-232,632. Examples of reverse transcriptase inhibitors include delaviridine, efavirenz, GS-840, HB Y097, lamivudine, nevirapine, AZT, 3TC, ddC, and ddI.

“Angiogenesis inhibitors” refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR2), inhibitors of epidermal-derived, fibroblast-derived, or platelet derived growth factors, MMP (matrix metalloprotease) inhibitors, integrin blockers, interferon- α , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (PNAS, Vol. 89, p. 7384 (1992); JNCI, Vol. 69, p. 475 (1982); Arch. Ophthalmol., Vol. 108, p.573 (1990); Anat. Rec.,

Vol. 238, p. 68 (1994); FEBS Letters, Vol. 372, p. 83 (1995); Clin. Orthop. Vol. 313, p. 76 (1995); J. Mol. Endocrinol., Vol. 16, p.107 (1996); Jpn. J. Pharmacol., Vol. 75, p. 105 (1997); Cancer Res., Vol. 57, p. 1625 (1997); Cell, Vol. 93, p. 705 (1998); Intl. J. Mol. Med., Vol. 2, p. 715 (1998); J. Biol. Chem., Vol. 274, p. 9116 (1999)),
5 carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, angiotensin II antagonists (see Fernandez et al., J. Lab. Clin. Med. 105:141-145 (1985)), and antibodies to VEGF (see, Nature Biotechnology, Vol. 17, pp.963-968 (October 1999); Kim et al., Nature, 362, 841-844 (1993); WO 00/44777; and WO 00/61186).

10 As described above, the combinations with NSAID's are directed to the use of NSAID's which are potent COX-2 inhibiting agents. For purposes of this specification an NSAID is potent if it possess an IC₅₀ for the inhibition of COX-2 of 1 μ M or less as measured by the cell or microsomal assay disclosed herein.

The invention also encompasses combinations with NSAID's which
15 are selective COX-2 inhibitors. For purposes of this specification NSAID's which are selective inhibitors of COX-2 are defined as those which possess a specificity for inhibiting COX-2 over COX-1 of at least 100 fold as measured by the ratio of IC₅₀ for COX-2 over IC₅₀ for COX-1 evaluated by the cell or micromsal assay disclosed hereinunder. Such compounds include, but are not limited to those disclosed in
20 U.S. 5,474,995, issued December 12, 1995, U.S. 5,861,419, issued January 19, 1999, U.S. 6,001,843, issued December 14, 1999, U.S. 6,020,343, issued February 1, 2000, U.S. 5,409,944, issued April 25, 1995, U.S. 5,436,265, issued July 25, 1995, U.S. 5,536,752, issued July 16, 1996, U.S. 5,550,142, issued August 27, 1996, U.S. 5,604,260, issued February 18, 1997, U.S. 5,698,584, issued December 16, 1997,
25 U.S. 5,710,140, issued January 20, 1998, WO 94/15932, published July 21, 1994, U.S. 5,344,991, issued June 6, 1994, U.S. 5,134,142, issued July 28, 1992, U.S. 5,380,738, issued January 10, 1995, U.S. 5,393,790, issued February 20, 1995, U.S. 5,466,823, issued November 14, 1995, U.S. 5,633,272, issued May 27, 1997, and U.S. 5,932,598, issued August 3, 1999, all of which are hereby incorporated by
30 reference.

Other examples of specific inhibitors of COX-2 include the following:

- 3-(3-fluorophenyl)-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone;
- 3-(3,4-difluorophenyl)-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone;
- 3-(3,4-dichlorophenyl)-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone;
- 5 3-phenyl-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone;
- 5,5-dimethyl-3-(3-fluorophenyl)-4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone;
- 3-(4-methylsulfonyl)phenyl)-2-phenyl-5-trifluoromethylpyridine;
- 2-(3-chlorophenyl)-3-(4-methylsulfonyl)phenyl-5-trifluoromethyl-pyridine;
- 2-(4-chlorophenyl)-3-(4-methylsulfonyl)phenyl-5-trifluoromethyl-pyridine;
- 10 2-(4-fluorophenyl)-3-(4-methylsulfonyl)phenyl-5-trifluoromethyl-pyridine;
- 3-(4-methylsulfonyl)phenyl)-2-(3-pyridinyl)-5-trifluoromethylpyridine;
- 5-methyl-3-(4-methylsulfonyl)phenyl)-2-phenylpyridine;
- 2-(4-chlorophenyl)-5-methyl-3-(4-methylsulfonyl) phenylpyridine;
- 5-methyl-3-(4-methylsulfonyl)phenyl)-2-(3-pyridinyl) pyridine;
- 15 5-chloro-2-(4-chlorophenyl)-3-(4-methylsulfonyl) phenylpyridine;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(2-pyridinyl) pyridine;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(3-pyridinyl) pyridine;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(4-pyridinyl) pyridine;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(2-methyl-5-pyridinyl)pyridine;
- 20 2-(4-chlorophenyl)-3-(4-methylsulfonyl)phenylpyridinyl-5-carboxylic acid methyl ester;
- 2-(4-chlorophenyl)-3-(4-methylsulfonyl)phenylpyridinyl-5-carboxylic acid;
- 5-cyano-2-(4-chlorophenyl)-3-(4-methylsulfonyl) phenylpyridine;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(3-pyridyl)pyridine hydromethanesulfonate;
- 25 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(3-pyridyl)pyridine hydrochloride;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(2-methyl-5-pyridinyl)pyridine hydrochloride;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(2-ethyl-5-pyridinyl)pyridine;
- 5-chloro-3-(4-methylsulfonyl)phenyl)-2-(2-ethyl-5-pyridinyl)pyridine hydromethanesulfonate;
- 30 3-(3,4-difluorophenoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5*H*-furan-2-one;

- 3-(3-fluorophenoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(3,5-difluorophenoxy)-5,5-dimethyl-4-(methylsulfonyl) phenyl)-5H-furan-2-one;
3-phenoxy-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(2,4-difluorophenoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
5 3-(4-chlorophenoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(3,4-dichlorophenoxy)-5,5-dimethyl-4-(methylsulfonyl) phenyl)-5H-furan-2-one;
3-(4-fluorophenoxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
3-(4-fluorophenylthio)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(3,5-difluorophenylthio)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-
10 one;
3-phenylthio-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(N-phenylamino)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
3-(N-methyl-N-phenylamino)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-
one;
15 3-cyclohexyloxy-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-phenylthio-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-benzyl-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(3,4-difluorophenylhydroxymethyl)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-
furan-2-one;
20 3-(3,4-difluorobenzoyl)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-benzoyl-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
4-(4-(methylsulfonyl)phenyl)-3-phenoxy-1-oxaspiro[4.4]non-3-en- 2-one;
4-(4-(methylsulfonyl)phenyl)-3-phenylthio-1-oxaspiro[4.4]non-3-en-2-one;
4-(2-oxo-3-phenylthio-1-oxa-spiro[4.4]non-3-en-4-yl) benzenesulfonamide;
25 3-(4-fluorobenzyl)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
3-(3,4-difluorophenoxy)-5-methoxy-5-methyl-4-(4- (methylsulfonyl)phenyl)-5H-
furan-2-one;
3-(5-chloro-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-
one;
30 3-(2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;

- 3-(6-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
- 3-(3-isoquinolinoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(4-(methylsulfonyl)phenyl)-2-phenoxy-cyclopent-2-enone;
- 5 3-(4-(methylsulfonyl)phenyl)-2-(3,4-difluorophenoxy)cyclopent-2-enone;
- 5,5-dimethyl-4-(4-methylsulfonylphenyl)-3-(5-bromopyridin-2-yloxy)-5H-furan-2-one;
- 5,5-dimethyl-4-(4-methylsulfonylphenyl)-3-(2-propoxy)-5H-furan-2-one;
- 2-(3,4-difluorophenoxy)-3-(4-methylsulfonylphenyl)-cyclopent-2-enone;
- 10 3-(5-benzothiophenyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
- 5,5-dimethyl-4-(4-methylsulfonyl-phenyl)-3-(pyridyl-4-oxy)-5H-furan-2-one;
- 5,5-dimethyl-4-(4-methylsulfonyl-phenyl)-3-(pyridyl-3-oxy)-5H-furan-2-one;
- 3-(2-methyl-5-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-
- 15 one;
- 3-(2-fluoro-4-trifluoromethyl)phenoxy-4-(4-methylsulfonyl)phenyl)-5,5-dimethyl-5H-furan-2-one;
- 3-(5-chloro-2-pyridylthio)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 2-(3,5-difluorophenoxy)-3-(4-methylsulfonylphenyl)-cyclopent-2-enone;
- 20 3-(2-pyrimidinoxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 3-(3-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 3-(3-chloro-5-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
- 3-(3-(1,2,5-thiadiazolyl)oxy)-4-(4-(methylsulfonyl)phenyl)-5,5-dimethyl-5H-furan-2-
- 25 one;
- 3-(5-isoquinolinoxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 3-(6-amino-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
- 3-(3-chloro-4-fluoro)phenoxy-4-(methylsulfonyl)phenyl)-5,5-dimethyl-5H-furan-2-
- 30 one;

- 3-(6-quinolinoxyl)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(5-nitro-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(2-thiazolylthio)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(3-chloro-5-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-
5 one;
5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-3-(2-propoxy)-5H-furan-2-one;
3-(3-trifluoromethyl)phenoxy-4-(4-(methylsulfonyl)phenyl)-5,5-dimethyl-5H-furan-2-
one;
5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-3-(piperidine-1-carbonyl)-5H-furan-2-
10 one;
5,5-dimethyl-3-(2-butoxy)-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-3-(3-pentoxy)-5H-furan-2-one;
2-(5-chloro-2-pyridyloxy)-3-(4-(methylsulfonyl)phenyl)cyclopent-2-enone;
3-(4-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
15 (5R)-3-(3,4-difluorophenoxy)-5-ethyl-5-methyl-4-(4-(methylsulfonyl)phenyl)-5H-
furan-2-one;
(5R)-3-(4-chlorophenoxy)-5-ethyl-5-methyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-
one;
3-(2-methyl-3-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
20 3-(4-methyl-5-nitro-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-
furan-2-one;
3-(5-chloro-4-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-
furan-2-one;
3-(5-fluoro-4-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-
25 furan-2-one;
3-(3-chloro-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
3-(4-fluorophenoxy)-5-methyl-4-(4-(methylsulfonyl)phenyl)-5-propyl-5H-furan-2-one;
3-(N,N-diethylamino)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-3-(3,5-dichloro-2-pyridyloxy)-5H-furan-2-
30 one;

- (5R)-3-(4-bromophenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(4-methoxyphenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 5 (5R)-3-(5-chloro-2-pyridyloxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5-(2,2,2-trifluoroethyl)-5H-furan-2-one;
- 3-(5-chloro-2-pyridyloxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5-propyl-5H-furan-2-one;
- 3-(1-cyclopropyl-ethoxy)-5,5-dimethyl-4-(4-methyl sulfonyl)phenyl)-5H-furan-2-one;
- 10 5-methyl-4-(4-(methylsulfonyl)phenyl)-3-(2-(propoxy)-5-(2-trifluoroethyl)-5H-furan-2-one;
- 5(R)-5-ethyl-5-methyl-4-(4-(methylsulfonyl)phenyl)-3-(2-propoxy)-5H-furan-2-one;
- 5,5-dimethyl-3-(2,2-dimethylpropyloxy)-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 15 5(R)-3-(1-cyclopropyl-ethoxy)-5-ethyl-5-methyl-4-(4-(methyl sulfonyl)phenyl)-5H-furan-2-one;
- 5(S)-5-ethyl-5-methyl-4-(4-(methylsulfonyl)phenyl)-3-(2-propoxy)-5H-furan-2-one;
- 3-(1-cyclopropylethoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(1-cyclopropylethoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 20 5,5-dimethyl-3-(isobutoxy)-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(4-bromophenoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(2-quinolinoxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(2-chloro-5-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 25 3-(6-benzothiazolyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
- 3-(6-chloro-2-pyridyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl) phenyl)-5H-furan-2-one;
- 3-(4-quinazolyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- (5R)-3-(5-fluoro-2-pyridyloxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 30 furan-2-one;


- (5R)-3-(4-fluorophenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(5-fluoro-2-pyridyloxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5-(2,2,2-trifluoroethyl)-5H-furan-2-one;
- 5 3-(1-isoquinolinylxy)-5,5-dimethyl-4-(methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(4-fluorophenoxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5-(2,2,2-trifluoroethyl)-5H-furan-2-one;
- 3-(3-fluoro-2-pyridyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(3,4-difluorophenoxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5-(2,2,2-trifluoroethyl)-5H-furan-2-one;
- 10 (5R)-3-(5-chloro-2-pyridyloxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 3-(3,4-difluorophenoxy)-5-methyl-5-trifluoromethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 15 3-(3,4-difluorophenoxy)-5-methyl-4-(4-(methylsulfonyl)phenyl)-5-propyl-5H-furan-2-one;
- 3-cyclobutyloxy-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 3-(1-indanyloxy)-5,5-dimethyl-4-(4-(methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(2-indanyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl)-5H-furan-2-one;
- 20 3-cyclopentyloxy-5,5-dimethyl-4-(4-methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-(3,3-dimethylcyclopentyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl)-5H-furan-2-one;
- 3-isopropoxy-5-methyl-4-(4-methylsulfonyl)phenyl)-5-propyl-5H-furan-2-one;
- 3-(2-methoxy-5-pyridyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 25 one;
- 3-(5-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5RS)-3-(3,4-difluorophenoxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5-(2,2,2-trifluoroethyl)-5H-furan-2-one;
- 3-(3-chloro-4-methoxyphenoxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-
- 30 2-one;

- (5R)-3-(3-chloro-4-methoxyphenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(4-chlorophenoxy)-5-trifluoroethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 5 (5R)-3-(4-bromophenoxy)-5-trifluoroethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 5-cyclopropylmethyl-3-(3,4-difluorophenoxy)-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(3-fluorophenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-
- 10 one;
- (5R)-3-(4-chloro-3-fluorophenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-phenoxy-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(4-chloro-3-methylphenoxy)-5-ethyl-5-methyl-4-(4-methylsulfonyl)phenyl-
- 15 5H-furan-2-one;
- 3-(4-chloro-3-methylphenoxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- (5R)-3-(5-bromo-2-pyridyloxy)-4-(4-methylsulfonylphenyl)-5-methyl-5-(2,2,2-trifluoroethyl)-5H-furan-2-one;
- 20 (5R)-3-(5-bromo-2-pyridyloxy)-4-(4-methylsulfonylphenyl)-5-ethyl-5-methyl-5H-furan-2-one;
- 3-(5-chloro-6-methyl-2-pyridyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- 3-(5-cyclopropyl-2-pyridyloxy)-5,5-dimethyl-4-(4-methylsulfonyl)phenyl-5H-furan-2-
- 25 one;
- 3-(1-cyclopropylethoxy)-4-(4-methylsulfonyl)phenyl-5H-furan-2-one; and
- 3-(cyclopropylmethoxy)-4-(4-methylsulfonyl)phenyl-5H-furan-2-one;
- or a pharmaceutically acceptable salt or stereoisomer thereof.

Inhibitors of COX-2 that are particularly useful in the instant method of treatment are:


3-phenyl-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone; and

5



5-chloro-3-(4-methylsulfonyl)phenyl-2-(2-methyl-5-pyridinyl)pyridine;


10



or a pharmaceutically acceptable salt thereof.

General and specific synthetic procedures for the preparation of the COX-2 inhibitor compounds described above are found in U.S. Patent No. 5,474,995, issued December 12, 1995, U.S. Patent No. 5,861,419, issued January 19, 1999, and U.S. Patent No. 6,001,843, issued December 14, 1999, all of which are herein incorporated by reference.

Compounds that have been described as specific inhibitors of COX-2 and are therefore useful in the present invention include, but are not limited to, the following:



5 or a pharmaceutically acceptable salt thereof.

Compounds which are described as specific inhibitors of COX-2 and are therefore useful in the present invention, and methods of synthesis thereof, can be found in the following patents, pending applications and publications, which are herein incorporated by reference: WO 94/15932, published July 21, 1994, U.S. Patent No. 5,344,991, issued June 6, 1994, U.S. Patent No. 5,134,142, issued July 28, 1992, U.S. Patent No. 5,380,738, issued January 10, 1995, U.S. Patent No. 5,393,790, issued February 20, 1995, U.S. Patent No. 5,466,823, issued November 14, 1995, U.S. Patent No. 5,633,272, issued May 27, 1997, and U.S. Patent No. 5,932,598, issued August 3, 1999.

10

Compounds which are specific inhibitors of COX-2 and are therefore useful in the present invention, and methods of synthesis thereof, can be found in the following patents, pending applications and publications, which are herein incorporated by reference: U.S. Patent No. 5,474,995, issued December 12, 1995 ,
5 U.S. Patent No. 5,861,419, issued January 19, 1999, U.S. Patent No. 6,001,843, issued December 14, 1999, U.S. Patent No. 6,020,343, issued February 1, 2000, U.S. Patent No. 5,409,944, issued April 25, 1995, U.S. Patent No. 5,436,265, issued July 25, 1995, U.S. Patent No. 5,536,752, issued July 16, 1996, U.S. Patent No. 5,550,142, issued August 27, 1996, U.S. Patent No. 5,604,260, issued February 18,
10 1997, U.S. Patent No. 5,698,584, issued December 16, 1997, and U.S. Patent No. 5,710,140, issued January 20, 1998.

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin, ukrain, ranpirnase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-butenyl)oxiranyl]-1-oxaspiro[2,5]oct-6-yl(chloroacetyl)carbamate, acetyldinanaline,
15 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1H-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentaose phosphate, 7,7-(carbonyl-bis[imino-N-methyl-4,2-pyrrolocarbonyl-imino[N-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene disulfonate), and 3-[(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

20 As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_3$ integrin, to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_5$ integrin, to compounds which antagonize, inhibit or counteract binding of a physiological ligand to both the $\alpha_v\beta_3$
25 integrin and the $\alpha_v\beta_5$ integrin, and to compounds which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins. The term also refers to antagonists of any combination of $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins.

Some specific examples of tyrosine kinase inhibitors include N-(trifluoromethylphenyl)-5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one, SH268, genistein, STI571, CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7H-pyrrolo[2,3-d]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, SU6668, STI571A, N-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

The instant compounds are also useful, alone or in combination with platelet fibrinogen receptor (GP IIb/IIIa) antagonists, such as tirofiban, to inhibit metastasis of cancerous cells. Tumor cells can activate platelets largely via thrombin generation. This activation is associated with the release of VEGF. The release of VEGF enhances metastasis by increasing extravasation at points of adhesion to vascular endothelium (Amirkhosravi, *Platelets* 10, 285-292, 1999). Therefore, the present compounds can serve to inhibit metastasis, alone or in combination with GP IIb/IIIa) antagonists. Examples of other fibrinogen receptor antagonists include abciximab, eptifibatide, sibrifiban, lamifiban, lotrafiban, cromofiban, and CT50352.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other pharmaceutically active agent(s) within its approved dosage range. Compounds of the instant invention may alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a combination formulation is inappropriate.

The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in

combination with one or more other active agents (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

The present invention also encompasses a pharmaceutical composition useful in the treatment of cancer, comprising the administration of a therapeutically effective amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents. Suitable compositions of this invention include aqueous solutions comprising compounds of this invention and pharmacologically acceptable carriers, e.g., saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's bloodstream by local bolus injection.


When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg

of body weight per day.


The compounds of this invention may be prepared by employing reactions as shown in the following schemes, in addition to other standard manipulations that are known in the literature or exemplified in the experimental procedures. These schemes, therefore, are not limited by the compounds listed nor by any particular substituents employed for illustrative purposes. Substituent numbering as shown in the schemes do not necessarily correlate to that used in the claims.

SCHEME A




10

SCHEME A (CONTINUED)




SCHEME B

EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention and not limiting of the reasonable scope thereof.

SCHEME 1



N-methoxy-N-methyl-2-(methylthio)pyrimidine-4-carboxamide (1-2)

A mixture of 2-(methylthio)pyrimidine-4-carboxylic acid (1-1, prepared according to Kim, C.-H., *et al J. Med. Chem.* **1986**, 29, 1374-1380 and McOmie, W. *J. Chem. Soc.* **1953**, 3129, 2.00 g, 11.8 mmol, 1 equiv), *N,O*-dimethylhydroxylamine hydrochloride (3.44 g, 35.3 mmol, 3.00 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (3.38 g, 17.6 mmol, 1.50 equiv), 1-hydroxy-7-azabenzotriazole (2.40 g, 17.6 mmol, 1.50 equiv), and triethylamine (6.55 mL, 47.0 mmol, 4.00 equiv) in DMF (40 mL) was stirred for 20 hours. The reaction mixture was partitioned between a 1:1 mixture of brine and saturated aqueous sodium carbonate solution (100 mL) and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexane initially, grading to 40% hexane in ethyl acetate) to give N-methoxy-N-methyl-2-(methylthio)pyrimidine-4-carboxamide (1-2) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 8.63 (d, 1H, *J* = 4.9 Hz), 7.13 (br s, 1H), 3.74 (br s, 3H), 3.36 (br s, 3H), 2.59 (s, 3H).

1-[2-(methylthio)pyrimidin-4-yl]ethanone (1-3)

A solution of methylmagnesium bromide in diethyl ether (3.0 M, 6.10 mL, 18.3 mmol, 3.0 equiv) was added to a solution of N-methoxy-N-methyl-2-(methylthio)pyrimidine-4-carboxamide (1-2, 1.30 g, 6.10 mmol, 1 equiv) in THF at -78°C. The reaction mixture was warmed to 0°C, stirred for 30 minutes, then partitioned between brine and ethyl acetate (2 x 75 mL). The combined organic layers were dried over sodium sulfate and concentrated to give 1-[2-(methylthio)pyrimidin-4-yl]ethanone (1-3) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 8.74 (d, 1H, *J* = 4.9 Hz), 7.51 (d, 1H, *J* = 4.9 Hz), 2.70 (s, 3H), 2.63 (s, 3H).

2-bromo-1-[2-(methylthio)pyrimidin-4-yl]ethanone (1-4)

A solution of 1-[2-(methylthio)pyrimidin-4-yl]ethanone (1-3, 906 mg, 5.4 mmol, 1 equiv) in 48% HBr (32 mL) was treated with bromine (860 mg, 5.4 mmol, 1 equiv) and stirred under ambient conditions for 18 hours. The insoluble yellow

solid was collected by filtration and partitioned between EtOAc and sat. NaHCO₃ solution. The organic layer was washed with brine, dried over MgSO₄ and concentrated to give 2-bromo-1-[2-(methylthio)pyrimidin-4-yl]ethanone (**1-4**) as a yellow solid. ¹H NMR (500 MHz, CDCl₃) δ 8.79 (d, 1H, *J* = 4.9 Hz), 7.56 (d, 1H, *J* = 4.9 Hz), 4.74 (s, 2H), 2.62 (s, 3H).

4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine hydrobromide (**1-5**)

A solution of 2-bromo-1-[2-(methylthio)pyrimidin-4-yl]ethanone (**1-4**, 700 mg, 2.8 mmol, 1 equiv), thiourea (215 mg, 2.8 mmol, 1 equiv) in ethanol (50 mL) was heated at 70°C for 30 minutes. The reaction mixture was cooled to provide 4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine hydrobromide (**1-5**) as a solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.64 (d, 1H, *J* = 5.1 Hz), 7.63 (br s, 1H), 7.48 (d, 1H, *J* = 5.1 Hz), 2.55 (s, 3H).

5-bromo-4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazol-2-amine (**1-6**)

A solution of oxone (367 mg, 0.60 mmol, 1.20 equiv) in water (5 mL) was added to a solution of 4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine hydrobromide (**1-5**, 152 mg, 0.50 mmol, 1 equiv) in methanol/water (1/1, 20 mL) at 23°C, and the resulting mixture was stirred for 2 hours. The reaction mixture made basic with solid NaHCO₃ and concentrated to remove the methanol. The aqueous residue was extracted with EtOAc (3 x 100 mL). The combined organic layers were dried over MgSO₄ and concentrated to give 5-bromo-4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazol-2-amine (**1-6**). ¹H NMR (500 MHz, CDCl₃) δ 8.86 (d, 1H, *J* = 5.1 Hz), 8.11 (d, 1H, *J* = 5.1 Hz), 6.78 (br s, 2H), 3.03 (s, 3H).


4-(2-amino-5-bromo-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine (**1-7**)

A mixture of 5-bromo-4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazol-2-amine (**1-6**, 130 mg, 0.41 mmol) and 3,5-dimethylaniline (0.5 mL, excess) was heated at 80°C for 18 hours. The residue was purified by flash column chromatography.

Elution with 20% EtOAc/hexanes to 80% EtOAc/hexanes gave 4-(2-amino-5-bromo-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine (**1-7**) as an off-white solid. ^1H NMR (500 MHz, CDCl_3) δ 8.48 (d, 1H, $J = 4.9$ Hz), 7.32 (m, 3H), 7.12 (s, 1H), 6.70 (s, 1H), 5.00 (s, 2H), 2.33 (s, 6H).

5

SCHEME 2



10 4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazol-2-amine (**2-1**)


4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine hydrobromide (**1-5**, 152 mg, 0.5 mmol, 1 equiv) was sonicated with sat. NaHCO_3 solution to give 4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine which was dissolved in methanol (30 mL). A solution of oxone (368 mg, 0.6 mmol, 1.2 equiv) in water (5 mL) was added to the methanol solution and the reaction was stirred under ambient conditions for 18 hours. The reaction was concentrated and the residue was partitioned between EtOAc and sat. NaHCO_3 solution. The organic layer was dried over MgSO_4 and concentrated to give 4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazol-2-amine (**2-1**)

as an off-white solid. ^1H NMR (500 MHz, CDCl_3) δ 8.87 (d, 1H, $J = 5.1$ Hz), 8.00 (d, 1H, $J = 5.1$ Hz), 7.83 (s, 1H), 5.10 (br s, 2H), 3.39 (s, 3H).

4-(2-amino-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine (2-2)

5 A mixture of 4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazol-2-amine (2-1, 114 mg, 0.45 mmol) and 3,5-dimethylaniline (1 mL, excess) was heated at 80°C for 18 hours. The residue was purified by chromatography. Elution with 20% EtOAc/hexanes to 80 % EtOAc/hexanes gave 4-(2-amino-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine (2-2) as an off-white solid. ^1H NMR (500 MHz, 10 CDCl_3) δ 8.46 (d, 1H, $J = 5.1$ Hz), 7.48 (s, 1H), 7.26 (s, 2H), 7.23 (d, 1H, $J = 5.1$ Hz), 7.07 (s, 1H), 6.70 (s, 1H), 5.01 (s, 2H), 2.34 (s, 6 H).

SCHEME 3



5 5-bromo-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine (3-1)

- 4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine (1-5, 466 mg, 1.53 mmol, 1 equiv) was suspended in 1/1 solution of 48% HBr and acetic acid (40 mL). This suspension was diluted with water to form a solution and then treated with bromine (244 mg, 1.53 mmol, 1 equiv). The reaction was stirred for 2 hours and then concentrated. The residue was partitioned between EtOAc and sat. NaHCO_3 solution. The organic layer was washed with brine, dried over MgSO_4 and concentrated to give 5-bromo-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine (3-1) as a pale yellow solid. ^1H NMR (500 MHz, CDCl_3) δ 8.55 (d, 1H, $J = 5.1$ Hz), 7.54 (d, 1H, $J = 5.1$ Hz), 5.00 (s, 2H), 2.67 (s, 3H).

15

4-[2-(methylthio)pyrimidin-4-yl]-5-phenyl-1,3-thiazol-2-amine (3-2)

The deoxygenated mixture of 5-bromo-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine (3-1, 410 mg, 1.35 mmol, 1 equiv), phenylboronic acid (165 mg, 1.53 mmol, 1 equiv), lithium chloride (172 mg, 3 mmol, 3 equiv), 2 M NaCO₃ (6.76 mL, 10 equiv) and tetrakis(triphenylphosphine)palladium(0) (78 mg, .07 mmol, 0.05 equiv) was heated under nitrogen at 90°C for 18 hours. The reaction was concentrated and the residue was partitioned between EtOAc and sat. NaHCO₃ solution. The organic layer was washed with brine, dried over MgSO₄ and concentrated to give a red gum which was purified by chromatography. Elution with 10% EtOAc/CH₂Cl₂ to 40% EtOAc/CH₂Cl₂ gave 4-[2-(methylthio)pyrimidin-4-yl]-5-phenyl-1,3-thiazol-2-amine (3-2) as a tan solid. ¹H NMR (500 MHz, CDCl₃) δ 8.44 (d, 1H, *J* = 5.1 Hz), 7.34 (m, 6 Hz), 4.99 (s, 2H), 1.93 (s, 3H).

4-[2-(methylsulfonyl)pyrimidin-4-yl]-5-phenyl-1,3-thiazol-2-amine (3-3)

An aqueous solution (5 mL) of oxone (166 mg, 0.27 mmol, 1.2 equiv) was added to a methanol solution (15 mL) of 4-[2-(methylthio)pyrimidin-4-yl]-5-phenyl-1,3-thiazol-2-amine (3-2, 68 mg, 0.23 mmol, 1 equiv) and the reaction was stirred under ambient conditions for 18 hours. The reaction was concentrated and the residue partitioned between EtOAc and sat. NaHCO₃ solution. The organic layer was washed with brine, dried over MgSO₄ and concentrated to provide 4-[2-(methylsulfonyl)pyrimidin-4-yl]-5-phenyl-1,3-thiazol-2-amine (3-3) as an off-white solid. ¹H NMR (500 MHz, CDCl₃) δ 8.86 (d, 1H, *J* = 5.1 Hz), 8.81 (d, 1H, *J* = 5.1 Hz), 7.41 (m, 5 Hz), 5.15 (s, 2H), 2.61 (s, 3H).


4-(2-amino-5-phenyl-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine (3-4)

A mixture of 4-[2-(methylsulfonyl)pyrimidin-4-yl]-5-phenyl-1,3-thiazol-2-amine (3-3, 74 mg, 0.22 mmol), and 3,5-dimethylaniline (1 mL, excess) was heated at 90°C for 18 hours. The residue was purified by chromatography. Elution with 20% EtOAc/hexanes to EtOAc gave 4-(2-amino-5-phenyl-1,3-thiazol-4-yl)-N-

(3,5-dimethylphenyl)pyrimidin-2-amine (**3-4**) as a tan foam. ^1H NMR (500 MHz, CDCl_3) δ 8.28 (d, 1H, $J = 5.1$ Hz), 7.38 (m, 2H), 7.32 (m, 3 Hz), 7.25 (m, 2H), 7.12 (s, 2H), 7.05 (br s, 1H), 6.70 (m, 1H), 6.65 (s, 1H), 5.03 (s, 2H), 2.27 (s, 6H)

5

SCHEME 4



2-amino-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-1**)

10

A deoxygenated DMF mixture (10 mL) of 5-bromo-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine (**3-1**, 93 mg, 0.37 mmol, 1 equiv), zinc cyanide (22 mg, 0.18 mmol, 0.6 equiv), 1,1'-bis(diphenylphosphino)ferrocene (41 mg, 0.07 mmol, 0.24 equiv) and tris(dibenzylideneacetone) dipalladium(0) (28 mg, 0.03

mmol, 0.1 equiv) was placed in a 120°C oil bath for 30 minutes. The reaction was concentrated and the residue was partitioned between EtOAc and sat. NaHCO₃ solution. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was triturated with CH₂Cl₂ to provide 2-amino-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-1**) as a brown solid. ¹H NMR (500 MHz, CD₃OD) δ 8.63 (d, 1H, *J* = 5.1 Hz), 7.64 (d, 1H, *J* = 5.1 Hz), 2.68 (s, 3H).

2-amino-4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-2**) and 2-amino-4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-3**)

10 Aqueous solution (5 ml) of oxone (169 mg, 0.27 mmol, 1.2 equiv) was added to a methanol solution (100 mL) of 2-amino-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-1**, 57 mg, 0.23 mmol, 1 equiv) and the reaction was stirred under ambient conditions for 18 hours. The reaction was concentrated and the residue was partitioned between EtOAc and sat. NaHCO₃ solution. The organic
15 layer was washed with brine, dried over MgSO₄ and concentrated to give a mixture (3/2) of 2-amino-4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-2**) and 2-amino-4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-3**), respectively.

Calc'd for C₉H₇N₅O₂S₂ (M + H) 282.0, found 282.0.

20 Calc'd for C₉H₇N₅OS₂ (M + H) 266.0, found 266.0.


2-amino-4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile (**4-4**)

A mixture (3/2) of 2-amino-4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-2**) and 2-amino-4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (**4-3**) (61 mg, 0.22 mmol) and 3,5-dimethylaniline (1 mL, excess) was heated at 90°C for 18 hours. The reaction was diluted with CH₂Cl₂ to give a yellow solid which was purified by reverse-phase LC (H₂O/CH₃CN gradient w/0.1% TFA present) to give 2-amino-4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-

yl}-1,3-thiazole-5-carbonitrile (**4-4**). ^1H NMR (500 MHz, DMSO- d_6) δ 9.51 (s, 1H), 8.62 (d, 1H, $J = 4.9$ Hz), 8.28 (s, 2H), 7.47 (s, 2H), 7.21 (d, 1H, $J = 4.9$ Hz), 6.59 (s, 1H), 2.25 (s, 6H).

5

SCHEME 5



4-(5-bromo-1,3-thiazol-4-yl)-2-(methylthio)pyrimidine (**5-1**)

10

A cold aqueous solution (2.5 mL) of sodium nitrite (423 mg, 6.1 mmol, 1.2 equiv) was slowly added below the surface to a cooled (-5°C) 50% H_2SO_4 solution (50 mL) of 5-bromo-4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazol-2-amine (**3-1**, 1.55 g, 5.1 mmol, 1 equiv) so that the temperature did not rise above 0°C . This reaction was added to a 30% hypophosphorous acid solution (13 mL, excess) which was

15

cooled to -5°C at such a rate that the temperature did not exceed 0°C . After stirring

1 hour at 0°C, 10 N NaOH was added to adjust the pH to 6. Solid Na₂CO₃ was then added to adjust the pH to 8. This mixture was extracted with EtOAc (3 x 50 mL). The extracts were combined, dried over MgSO₄ and concentrated to provide 4-(5-bromo-1,3-thiazol-4-yl)-2-(methylthio)pyrimidine (5-1) as a tan solid. ¹H NMR (500 MHz, DMSO-d₆) δ 9.30 (s, 1H), 8.74 (d, 1H, *J* = 5.1 Hz), 7.76 (d, 1H, *J* = 5.1 Hz), 2.64 (s, 3H).

4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-2)

A deoxygenated DMF mixture (5 mL) of 4-(5-bromo-1,3-thiazol-4-yl)-2-(methylthio)pyrimidine (5-1, 1.2 g, 4.16 mmol, 1 equiv), zinc cyanide (293 mg, 2.5 mmol, 0.6 equiv), 1,1'-bis(diphenylphosphino)ferrocene (554 mg, 1.0 mmol, 0.24 equiv) and tris(dibenzylideneacetone) dipalladium(0) (381 mg, 0.42 mmol, 0.1 equiv) was placed in a 90°C oil bath for 60 minutes. The reaction was concentrated and the residue was partitioned between EtOAc and sat. NaHCO₃ solution. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was purified by chromatography. Elution with CH₂Cl₂ to 2% EtOAc/CH₂Cl₂ gave 4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-2) as a grey solid. ¹H NMR (500 MHz, CDCl₃) δ 9.02 (s, 1H), 8.70 (d, 1H, *J* = 5.1 Hz), 7.76 (d, 1H, *J* = 5.1 Hz), 2.75 (s, 3H).

4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-3) and

4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-4)

A aqueous solution (5 ml) of oxone (2.1g, 3.43 mmol, 1.2 equiv) was added to a methanol solution (250 mL) of 4-[2-(methylthio)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-2, 670 mg, 2.86 mmol, 1 equiv) and the reaction was stirred under ambient conditions for 18 hours. The reaction was concentrated and the residue sonicated with sat NaHCO₃ solution to give a mixture (3/7) of 4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-3) and 4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-4) respectively.


Calc'd for $C_9H_6N_4O_2S_2$ (M + H) 267.0, found 267.0.

Calc'd for $C_9H_6N_4OS_2$ (M + H) 250.0, found 250.0.





4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile (5-5)





- 5 A mixture(3/7) of 4-[2-(methylsulfonyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-3) and 4-[2-(methylsulfinyl)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile (5-4) (266 mg, 1.0 mmol) and 3,5-dimethylaniline(1 mL, excess) was heated at 90°C for 18 hours. The residue was purified by flash column chromatography. Elution with CH_2Cl_2 to 4% EtOAc/ CH_2Cl_2 give 4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-
- 10 1,3-thiazole-5-carbonitrile (5-5). 1H NMR (500 MHz, $CDCl_3$) δ 9.00 (s, 1H), 8.63 (d, 1H, $J = 5.1$ Hz), 7.50 (d, 1H, $J = 5.1$ Hz), 7.31 (s, 2H), 7.21 (s, 1H), 6.72 (s, 1H), 2.35 (s, 6H).

SCHEME 6







The following compounds are prepared using the general techniques illustrated and described above by replacing the aniline with a suitably substituted aniline:





6-1		4-[2-({3-[(4-acetylpiperazin-1-yl) methyl]-5-methylphenyl} amino) pyrimidin-4-yl]-2-amino-1,3-thiazole-5-carbonitrile
6-2		4-[2-({3-[(4-acetylpiperazin-1-yl) methyl]-5-chlorophenyl} amino)pyrimidin-4-yl]-2-amino-1,3-thiazole-5-carbonitrile
6-3		2-amino-4-{2-[(3-methyl-5-{[4-(methylsulfonyl) piperazin-1-yl]methyl} phenyl) amino] pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-4		2-amino-4-{2-[(3-chloro-5-{[4-(methylsulfonyl) piperazin-1-yl]methyl} phenyl) amino] pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile





6-5		2-amino-4-[2-((3-methyl-5-[(4-methylpiperazin-1-yl)carbonyl]phenyl)amino)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-6		2-amino-4-[2-((3-chloro-5-[(4-methylpiperazin-1-yl)carbonyl]phenyl)amino)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-7		2-amino-4-(2-([3-methyl-5-(piperazin-1-ylcarbonyl)phenyl]amino)pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile
6-8		2-amino-4-(2-([3-chloro-5-(piperazin-1-ylcarbonyl)phenyl]amino)pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile





6-9		2-amino-4-(2-{3-methyl-5-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile
6-10		2-amino-4-(2-{3-chloro-5-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile
6-11		2-amino-4-(2-{4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile
6-12		4-(3-{[4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}-5-methylbenzyl)-N-methylpiperazine-1-carboxamide






6-13		4-(3-{{4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl} amino}-5-chlorobenzyl)-N-methylpiperazine-1-carboxamide
6-14		4-(3-{{4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl} amino}-5-methylbenzyl)piperazine-1-carboxamide
6-15		4-(3-{{4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl} amino}-5-chlorobenzyl)piperazine-1-carboxamide
6-16		2-amino-4-{2-[(3-methyl-5-{4-(methylsulfonyl)piperidin-1-yl)methyl}phenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile






6-17		2-amino-4-{2-[(3-chloro-5-{[4-(methylsulfonyl)piperidin-1-yl]methyl}phenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-18		2-amino-4-[2-({3-methyl-5-[(4-methyl-5-oxo-1,4-diazepan-1-yl)methyl]phenyl}amino)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-19		2-amino-4-[2-({3-chloro-5-[(4-methyl-5-oxo-1,4-diazepan-1-yl)methyl]phenyl}amino)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-20		2-amino-4-(2-({4-(piperazin-1-ylcarbonyl)phenyl}amino)pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile






6-21		4-[2-({2-[(4-acetylpiperazin-1-yl)methyl]-6-methylpyridin-4-yl}amino)pyrimidin-4-yl]-2-amino-1,3-thiazole-5-carbonitrile
6-22		4-[2-({2-[(4-acetylpiperazin-1-yl)methyl]-6-chloropyridin-4-yl}amino)pyrimidin-4-yl]-2-amino-1,3-thiazole-5-carbonitrile
6-23		2-amino-4-{2-[(2-methyl-6-{[4-(methylsulfonyl)piperazin-1-yl]methyl}pyridin-4-yl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-24		2-amino-4-{2-[(2-chloro-6-{[4-(methylsulfonyl)piperazin-1-yl]methyl}pyridin-4-yl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile





6-25		4-[(4-{[4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}-6-methylpyridin-2-yl)methyl]-N-methylpiperazine-1-carboxamide
6-26		4-[(4-{[4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}-6-chloropyridin-2-yl)methyl]-N-methylpiperazine-1-carboxamide
6-27		4-[2-((3-((4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile)methyl)-5-methylphenyl)amino]piperazine-1-carboxamide
6-28		4-[2-((3-((4-(2-amino-5-cyano-1,3-thiazol-4-yl)pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile)methyl)-5-chlorophenyl)amino]piperazine-1-carboxamide


6-29		4-{2-[(3-methyl-5-[[4-(methylsulfonyl)piperazin-1-yl]methyl]phenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-30		4-{2-[(3-chloro-5-[[4-(methylsulfonyl)piperazin-1-yl]methyl]phenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-31		4-[2-({3-methyl-5-[(4-methylpiperazin-1-yl)carbonyl]phenyl)amino]pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-32		4-[2-({3-chloro-5-[(4-methylpiperazin-1-yl)carbonyl]phenyl)amino]pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile

6-33		4-[2-({3-methyl-5-[(piperazin-1-yl) carbonyl]phenyl}amino) pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-34		4-[2-({3-chloro-5-[(piperazin-1-yl) carbonyl]phenyl}amino) pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-35		4-(2-{3-methyl-5-[(4-methylpiperazin-1-yl) sulfonyl] phenyl} pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile
6-36		4-(2-{3-chloro-5-[(4-methylpiperazin-1-yl) sulfonyl] phenyl} pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile
6-37		4-(2-{4-[(4-methylpiperazin-1-yl) sulfonyl]phenyl}pyrimidin-4-yl)-1,3-thiazole-5-carbonitrile

6-38		4-(3-methyl-5-{{[4-(5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}benzyl)-N-methylpiperazine-1-carboxamide
6-39		4-(3-chloro-5-{{[4-(5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}benzyl)-N-methylpiperazine-1-carboxamide
6-40		4-(3-methyl-5-{{[4-(5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}benzyl)piperazine-1-carboxamide
6-41		4-(3-chloro-5-{{[4-(5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}benzyl)piperazine-1-carboxamide
6-42		4-{2-[(3-methyl-5-{{[4-(methylsulfonyl)piperidin-1-yl]methyl}phenyl]amino}pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile

6-43		4-{2-[(3-chloro-5- {[4-(methylsulfonyl) piperidin-1-yl]methyl} phenyl)amino]pyrimidin -4-yl}-1,3-thiazole-5- carbonitrile
6-44		4-[2-({3-methyl-5- [(4-methyl-5-oxo-1,4- diazepan-1-yl)methyl] phenyl}amino)pyrimidin -4-yl]-1,3-thiazole-5- carbonitrile
6-45		4-[2-({3-chloro-5- [(4-methyl-5-oxo-1,4- diazepan-1-yl)methyl] phenyl}amino)pyrimidin -4-yl]-1,3-thiazole-5- carbonitrile
6-46		4-(2-{[4-(piperazin-1- ylcarbonyl)phenyl] amino}pyrimidin-4- yl)-1,3-thiazole-5- carbonitrile
6-47		4-[2-({2-[(4- acetylpiperazin-1-yl) methyl]-6- methylpyridin-4-yl} amino)pyrimidin-4-yl]- 1,3-thiazole-5- carbonitrile

6-48		4-[2-({2-[(4-acetylpiperazin-1-yl) methyl]-6-chloropyridin-4-yl} amino)pyrimidin-4-yl]-1,3-thiazole-5-carbonitrile
6-49		4-{2-[(2-methyl-6-{{4-(methylsulfonyl) piperazin-1-yl} methyl} pyridin-4-yl) amino] pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-50		4-{2-[(2-chloro-6-{{4-(methylsulfonyl) piperazin-1-yl} methyl} pyridin-4-yl) amino] pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile
6-51		4-[(4-{{4-(5-cyano-1,3-thiazol-4-yl) pyrimidin-2-yl} amino}-6-methylpyridin-2-yl) methyl]-N-methylpiperazine-1-carboxamide

6-52		4-[(6-chloro-4-{[4-(5-cyano-1,3-thiazol-4-yl)pyrimidin-2-yl]amino}pyridin-2-yl)methyl]-N-methylpiperazine-1-carboxamide
------	--	---

ASSAYS

- 5 The compounds of the instant invention described in the Examples were tested by the assays described below and were found to have kinase inhibitory activity. Other assays are known in the literature and could be readily performed by those of skill in the art. (see, for example, Dhanabal et al., *Cancer Res.* 59:189-197; Xin et al., *J. Biol. Chem.* 274:9116-9121; Sheu et al., *Anticancer Res.* 18:4435-4441;
- 10 Ausprunk et al., *Dev. Biol.* 38:237-248; Gimbrone et al., *J. Natl. Cancer Inst.* 52:413-427; Nicosia et al., *In Vitro* 18:538-549).

VEGF/KDR RECEPTOR KINASE ASSAY

- 15 VEGF receptor kinase activity is measured by incorporation of radio-labeled phosphate into polyglutamic acid, tyrosine, 4:1 (pEY) substrate. The phosphorylated pEY product is trapped onto a filter membrane and the incorporation of radio-labeled phosphate quantified by scintillation counting.

MATERIALS

20

VEGF receptor kinase

The intracellular tyrosine kinase domains of human KDR (Terman, B.I. et al. *Oncogene* (1991) vol. 6, pp. 1677-1683.) and Flt-1 (Shibuya, M. et al. *Oncogene* (1990) vol. 5, pp. 519-524) were cloned as glutathione S-transferase

(GST) gene fusion proteins. This was accomplished by cloning the cytoplasmic domain of the KDR kinase as an in frame fusion at the carboxy terminus of the GST gene. Soluble recombinant GST-kinase domain fusion proteins were expressed in *Spodoptera frugiperda* (Sf21) insect cells (Invitrogen) using a baculovirus expression
5 vector (pAcG2T, Pharmingen).

Lysis buffer

50 mM Tris pH 7.4, 0.5 M NaCl, 5 mM DTT, 1 mM EDTA, 0.5% triton X-100, 10% glycerol, 10 mg/mL of each leupeptin, pepstatin and aprotinin
10 and 1mM phenylmethylsulfonyl fluoride (all Sigma).

Wash buffer

50 mM Tris pH 7.4, 0.5 M NaCl, 5 mM DTT, 1 mM EDTA, 0.05% triton X-100, 10% glycerol, 10 mg/mL of each leupeptin, pepstatin and aprotinin and
15 1mM phenylmethylsulfonyl fluoride.

Dialysis buffer

50 mM Tris pH 7.4, 0.5 M NaCl, 5 mM DTT, 1 mM EDTA, 0.05% triton X-100, 50% glycerol, 10 mg/mL of each leupeptin, pepstatin and aprotinin and
20 1mM phenylmethylsulfonyl fluoride.

10 X reaction buffer

200 mM Tris, pH 7.4, 1.0 M NaCl, 50 mM MnCl₂, 10 mM DTT and 5 mg/mL bovine serum albumin (Sigma).
25

Enzyme dilution buffer

50 mM Tris, pH 7.4, 0.1 M NaCl, 1 mM DTT, 10% glycerol, 100 mg/mL BSA.

10 X Substrate

750 $\mu\text{g/mL}$ poly (glutamic acid, tyrosine; 4:1) (Sigma).

Stop solution

- 5 30% trichloroacetic acid, 0.2 M sodium pyrophosphate (both Fisher).

Wash solution

15% trichloroacetic acid, 0.2 M sodium pyrophosphate.

10 Filter plates

Millipore #MAFC NOB, GF/C glass fiber 96 well plate.

METHOD

15 A. Protein purification

1. Sf21 cells were infected with recombinant virus at a multiplicity of infection of 5 virus particles/ cell and grown at 27°C for 48 hours.
2. All steps were performed at 4°C. Infected cells were harvested by centrifugation at 1000 X g and lysed at 4°C for 30 minutes with 1/10 volume of
- 20 lysis buffer followed by centrifugation at 100,000 Xg for 1 hour. The supernatant was then passed over a glutathione Sepharose column (Pharmacia) equilibrated in lysis buffer and washed with 5 volumes of the same buffer followed by 5 volumes of wash buffer. Recombinant GST-KDR protein was eluted with wash buffer/10 mM reduced glutathione (Sigma) and dialyzed against dialysis buffer.

25

B. VEGF Receptor Kinase Assay

1. Add 5 μl of inhibitor or control to the assay in 50% DMSO.
2. Add 35 μl of reaction mix containing 5 μl of 10 X reaction buffer, 5 μl 25 mM ATP/10 μCi [^{33}P]ATP (Amersham), and 5 μl 10 X substrate.

3. Start the reaction by the addition of 10 μ l of KDR (25 nM) in enzyme dilution buffer.
4. Mix and incubate at room temperature for 15 minutes.
5. Stop by the addition of 50 μ l stop solution.
- 5 6. Incubate for 15 minutes at 4°C.
7. Transfer a 90 μ l aliquot to filter plate.
8. Aspirate and wash 3 times with wash solution.
9. Add 30 μ l of scintillation cocktail, seal plate and count in a Wallac Microbeta scintillation counter.

10

Human Umbilical Vein Endothelial Cell Mitogenesis Assay

- Expression of VEGF receptors that mediate mitogenic responses to the growth factor is largely restricted to vascular endothelial cells. Human umbilical vein endothelial cells (HUVECs) in culture proliferate in response to VEGF treatment and
- 15 can be used as an assay system to quantify the effects of KDR kinase inhibitors on VEGF stimulation. In the assay described, quiescent HUVEC monolayers are treated with vehicle or test compound 2 hours prior to addition of VEGF or basic fibroblast growth factor (bFGF). The mitogenic response to VEGF or bFGF is determined by measuring the incorporation of [³H]thymidine into cellular DNA.

20

Materials

HUVECs

- HUVECs frozen as primary culture isolates are obtained from
- 25 Clonetics Corp. Cells are maintained in Endothelial Growth Medium (EGM; Clonetics) and are used for mitogenic assays at passages 3-7.

Culture Plates

- NUNCCLON 96-well polystyrene tissue culture plates (NUNC
- 30 #167008).

Assay Medium

Dulbecco's modification of Eagle's medium containing 1 g/mL glucose (low-glucose DMEM; Mediatech) plus 10% (v/v) fetal bovine serum (Clonetics).

5

Test Compounds

Working stocks of test compounds are diluted serially in 100% dimethylsulfoxide (DMSO) to 400-fold greater than their desired final concentrations. Final dilutions to 1X concentration are made directly into Assay Medium immediately prior to addition to cells.

10

10X Growth factors

Solutions of human VEGF₁₆₅ (500 ng/mL; R&D Systems) and bFGF (10 ng/mL; R&D Systems) are prepared in Assay Medium.

15

10X [³H]Thymidine

[Methyl-³H]Thymidine (20 Ci/mmol; Dupont-NEN) is diluted to 80 µCi/mL in low-glucose DMEM.

20

Cell Wash Medium

Hank's balanced salt solution (Mediatech) containing 1 mg/mL bovine serum albumin (Boehringer-Mannheim).

Cell Lysis Solution

25

1 N NaOH, 2% (w/v) Na₂CO₃.

METHOD

1. HUVEC monolayers maintained in EGM are harvested by trypsinization and plated at a density of 4000 cells per 100 µL Assay Medium per

well in 96-well plates. Cells are growth-arrested for 24 hours at 37°C in a humidified atmosphere containing 5% CO₂.

2. Growth-arrest medium is replaced by 100 µL Assay Medium containing either vehicle (0.25% [v/v] DMSO) or the desired final concentration of test compound. All determinations are performed in triplicate. Cells are then
5 incubated at 37°C/5% CO₂ for 2 hours to allow test compounds to enter cells.

3. After the 2-hour pretreatment period, cells are stimulated by addition of 10 µL/well of either Assay Medium, 10X VEGF solution or 10X bFGF solution. Cells are then incubated at 37°C/5% CO₂.

10 4. After 24 hours in the presence of growth factors, 10X [³H]Thymidine (10 µL/well) is added.

5. Three days after addition of [³H]thymidine, medium is removed by aspiration, and cells are washed twice with Cell Wash Medium (400 µL/well followed by 200 µL/well). The washed, adherent cells are then solubilized
15 by addition of Cell Lysis Solution (100 µL/well) and warming to 37°C for 30 minutes. Cell lysates are transferred to 7-mL glass scintillation vials containing 150 µL of water. Scintillation cocktail (5 mL/vial) is added, and cell-associated radioactivity is determined by liquid scintillation spectroscopy.

Based upon the foregoing assays the compounds of Formula I are
20 inhibitors of VEGF and thus are useful for the inhibition of angiogenesis, such as in the treatment of ocular disease, e.g., diabetic retinopathy and in the treatment of cancers, e.g., solid tumors. The instant compounds inhibit VEGF-stimulated mitogenesis of human vascular endothelial cells in culture with IC₅₀ values between 0.01 - 5.0 µM. These compounds also show selectivity over related tyrosine kinases
25 (e.g., FGFR1 and the Src family; for relationship between Src kinases and VEGFR kinases, see Eliceiri et al., Molecular Cell, Vol. 4, pp.915-924, December 1999).

EREKA Kinase Assay (EGFR Assay)METHOD:

1. Dilute inhibitors (account for the final dilution into the assay, 1:20)
- 5 2. Prepare the appropriate amount of reaction mix at room temperature.
10X Buffer (20 mM Tris pH 7.4/0.1 M NaCl/1mM DTT final)
0.1M MnCl₂ (5mM final)
pEY substrate (75 µg/ml)
ATP/[³³P]ATP (2.5 µM/1 µCi final)
10 BSA (500 µg/ml final)
Na₃VO₄ (500 µM final)
3. Add 5 µl of the diluted inhibitor to the reaction mix. (Final volume of 5 µl in 50% DMSO) Positive control wells - add blank DMSO (50%).
4. Add 35 µl of the reaction mix to each well of a 96 well plate.
- 15 5. Dilute enzyme (1:100) into enzyme dilution buffer (keep at 4°C).
6. Add 10 µl of the diluted enzyme to each well and mix. Negative control wells - add 10 µl 0.5 M EDTA per well instead (final 100 mM)
7. Incubate at room temperature for 60 minutes.
8. Stop by the addition of an equal volume (50 µl) of 30% TCA/0.1M Na
20 pyrophosphate.
9. Incubate for 15 minutes to allow precipitation.
10. Transfer to Millipore filter plate.
11. Wash 3X with 15% TCA/0.1M Na pyrophosphate (125 µl per wash).
12. Allow to dry under vacuum for 2-3 minutes.
- 25 13. Dry in hood for ~ 20 minutes.
14. Assemble Wallac Millipore adapter and add 50 µl of scintillant to each well and count.

ALTERNATIVE METHOD:

- 30 1. Dilute inhibitors (account for the final dilution into the assay, 1:20)
2. Prepare the appropriate amount of reaction mix at room temperature.
10X Buffer (20 mM Tris pH 7.4/0.1 M NaCl/1mM DTT final)
0.1M MnCl₂ (5mM final)
pEY substrate (75 µg/ml)
35 ATP/[³³P]ATP (2.5 µM/1 µCi final)

BSA (500 $\mu\text{g/ml}$ final)

3. Add 5 μl of the diluted inhibitor to the reaction mix. (Final volume of 5 μl in 50% DMSO) Positive control wells - add blank DMSO (50%).
4. Add 35 μl of the reaction mix to each well of a 96 well plate.
- 5 5. Dilute enzyme into enzyme dilution buffer (keep at 4°C).
6. Add 10 μl of the diluted enzyme to each well and mix (5 nM final).
Negative control wells – add 10 μl 0.5 M EDTA per well instead (final 100 mM).
7. Incubate at room temperature for 30 minutes.
- 10 8. Stop by the addition of an equal volume (50 μl) of 30% TCA/0.1M Na pyrophosphate.
9. Incubate for 15 minutes to allow precipitation.
10. Transfer to Millipore filter plate.
11. Wash 3X with 15% TCA/0.1M Na pyrophosphate (125 μl per wash).
- 15 12. Allow to dry under vacuum for 2-3 minutes.
13. Dry in hood for ~ 20 minutes.
14. Assemble Wallac Millipore adapter and add 50 μl of scintillant to each well and count.

20

SRC ASSAYSRCKA (Mg⁺⁺) Kinase Assay

1. Dilute inhibitors (account for the final dilution into the assay, 1:20)
2. Prepare the appropriate amount of reaction mix at room temperature.
- 25 10X Buffer (20 mM Tris pH 7.4/0.1 M NaCl/1mM DTT final)
0.1M MgCl₂ (5mM final)
pEY substrate (75 $\mu\text{g/ml}$)
ATP/[³³P]ATP (24 μM /3 μCi final)
BSA (500 $\mu\text{g/ml}$ final)
- 30 3. Add 5 μl of the diluted inhibitor to the reaction mix. (Final volume of 5 μl in 50% DMSO)
Positive control wells - add blank DMSO (50%).
4. Add 35 μl of the reaction mix to each well of a 96 well plate.

5. Dilute enzyme (1:22) into enzyme dilution buffer (keep at 4°C). Final enzyme conc. = 0.44nM
6. Add 10 µl of the diluted enzyme to each well and mix.
Negative control wells – add 10 µl 0.5 M EDTA per well instead (final 100 mM)
- 5 7. Incubate at room temperature for 30 minutes.
8. Stop by the addition of an equal volume (50 µl) of 30% TCA/0.1M Na pyrophosphate.
9. Incubate for 15 minutes to allow precipitation.
10. Transfer to Millipore filter plate.
- 10 11. Wash 3X with 15% TCA/0.1M Na pyrophosphate (125 µl per wash).
12. Allow to dry under vacuum for 2-3 minutes.
13. Dry in hood for ~ 20 minutes.
14. Assemble Wallac Millipore adapter and add 50 µl of scintillant to each well and count.

15

ALTERNATIVE METHOD:

SRCKA (Mn++) Kinase Assay

1. Dilute inhibitors (account for the final dilution into the assay, 1:20).
2. Prepare the appropriate amount of reaction mix at room temperature.
- 20 10X Buffer (20 mM Tris pH 7.4/0.1 M NaCl/1mM DTT final)
0.1M MnCl₂ (5mM final)
pEY substrate (75 µg/ml)
ATP/[³³P]ATP (2.5 µM/1 µCi final)
BSA (500 µg/ml final)
- 25 3. Add 5 µl of the diluted inhibitor to the reaction mix. (Final volume of 5 µl in 50% DMSO) Positive control wells - add blank DMSO (50%).
4. Add 35 µl of the reaction mix to each well of a 96 well plate.
5. Dilute enzyme (1:44) into enzyme dilution buffer (keep at 4°C). Final enzyme conc. = 0.22nM
- 30 6. Add 10 µl of the diluted enzyme to each well and mix.
Negative control wells – add 10 µl 0.5 M EDTA per well instead (final 100 mM)
7. Incubate at room temperature for 30 minutes.
8. Stop by the addition of an equal volume (50 µl) of 30% TCA/0.1M Na pyrophosphate.

9. Incubate for 15 minutes to allow precipitation.
10. Transfer to Millipore filter plate.
11. Wash 3X with 15% TCA/0.1M Na pyrophosphate (125 µl per wash).
12. Allow to dry under vacuum for 2-3 minutes.
- 5 13. Dry in hood for ~ 20 minutes.
14. Assemble Wallac Millipore adapter and add 50 µl of scintillant to each well and count.


III. FLT-1 KINASE ASSAY

- 10 Flt-1 was expressed as a GST fusion to the Flt-1 kinase domain and was expressed in baculovirus/insect cells. The following protocol was employed to assay compounds for Flt-1 kinase inhibitory activity:
 1. Inhibitors were diluted to account for the final dilution in the assay, 1:20.
 - 15 2. The appropriate amount of reaction mix was prepared at room temperature:
 - 10X Buffer (20 mM Tris pH 7.4/0.1 M NaCl/1mM DTT final)
 - 0.1M MnCl₂ (5mM final)
 - pEY substrate (75 µg/mL)
 - ATP/[³³P]ATP (2.5 µM/1 µCi final)
 - 20 BSA (500 µg/mL final).
 3. 5 µL of the diluted inhibitor was added to the reaction mix. (Final volume of 5 µL in 50% DMSO). To the positive control wells, blank DMSO (50%) was added.
 4. 35 µL of the reaction mix was added to each well of a 96 well plate.
 - 25 5. Enzyme was diluted into enzyme dilution buffer (kept at 4°C).
 6. 10 µL of the diluted enzyme was added to each well and mix (5 nM final). To the negative control wells, 10 µL 0.5 M EDTA was added per well instead (final 100 mM).
 7. Incubation was then carried out at room temperature for 30 minutes.
 - 30 8. Stopped by the addition of an equal volume (50 µL) of 30% TCA/0.1M Na pyrophosphate.
 9. Incubation was then carried out for 15 minutes to allow precipitation.
 10. Transferred to Millipore filter plate.
 11. Washed 3X with 15% TCA/0.1M Na pyrophosphate (125 µL per wash).
 - 35 12. Allowed to dry under vacuum for 2-3 minutes.

13. Dried in hood for ~ 20 minutes.
14. Assembled Wallac Millipore adapter and added 50 μ L of scintillant to each well and counted.

WHAT IS CLAIMED IS:

1. A compound of Formula I



5

wherein

R^{1a} is independently selected from:

- (1) H,
 (2) unsubstituted or substituted C₁-C₁₀ alkyl,
 10 (3) OR⁸, and
 (4) N(R⁸)₂ ;

R¹ is independently selected from:

- (1) H,
 15 (2) unsubstituted or substituted C₁-C₁₀ alkyl,
 (3) unsubstituted or substituted C₃-C₁₀ cycloalkyl,
 (4) unsubstituted or substituted aryl,
 (5) unsubstituted or substituted heterocycle,
 (6) halo,
 20 (7) CF₃,
 (8) -(CH₂)_t R⁹C(O)R⁸,
 (9) -C(O)R⁹,
 (10) -(CH₂)_tOR⁸,
 (11) unsubstituted or substituted C₂-C₆ alkenyl,
 25 (12) unsubstituted or substituted C₂-C₆ alkynyl,
 (13) CN,

- (14) $-(\text{CH}_2)_t \text{NR}^7 \text{R}^8$,
- (15) $-(\text{CH}_2)_t \text{C}(\text{O})\text{NR}^7 \text{R}^8$,
- (16) $-\text{C}(\text{O})\text{OR}^8$, and
- (17) $-(\text{CH}_2)_t \text{S}(\text{O})_q (\text{CH}_2)_t \text{NR}^7 \text{R}^8$;

5

R^2 is independently selected from:

- (1) H,
- (2) unsubstituted or substituted $\text{C}_1\text{-C}_{10}$ alkyl,
- (3) unsubstituted or substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl,
- 10 (4) unsubstituted or substituted aryl,
- (5) unsubstituted or substituted heterocycle,
- (6) halo,
- (7) CF_3 ,
- (8) $-(\text{CH}_2)_t \text{R}^9 \text{C}(\text{O})\text{R}^8$,
- 15 (9) $-\text{C}(\text{O})\text{R}^9$,
- (10) $-(\text{CH}_2)_t \text{OR}^8$,
- (11) unsubstituted or substituted $\text{C}_2\text{-C}_6$ alkenyl,
- (12) unsubstituted or substituted $\text{C}_2\text{-C}_6$ alkynyl,
- (13) CN,
- 20 (14) $-(\text{CH}_2)_t \text{NR}^7 \text{R}^8$,
- (15) $-(\text{CH}_2)_t \text{C}(\text{O})\text{NR}^7 \text{R}^8$,
- (16) $-\text{C}(\text{O})\text{OR}^8$, and
- (17) $-(\text{CH}_2)_t \text{S}(\text{O})_q (\text{CH}_2)_t \text{NR}^7 \text{R}^8$;

25 R^3 is independently selected from:

- (1) H,
- (2) unsubstituted or substituted $\text{C}_1\text{-C}_{10}$ alkyl,
- (3) unsubstituted or substituted aralkyl,
- (4) CN,
- 30 (5) halo,
- (6) $\text{N}(\text{R}^8)_2$,
- (7) OR^8 , and
- (8) unsubstituted or substituted aryl;

R⁷ is selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
- (3) unsubstituted or substituted aralkyl;

5

R⁸ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- (3) unsubstituted or substituted aryl,
- 10 (4) unsubstituted or substituted heterocycle,
- (5) unsubstituted or substituted C₃-C₁₀ cycloalkyl, and
- (6) unsubstituted or substituted aralkyl;

15 R⁷ and R⁸, when attached to the same nitrogen atom may be joined to form a 5-7 membered heterocycle containing, in addition to the nitrogen, one or two more heteroatoms selected from N, O, or S, said heterocycle being optionally substituted with one to three R² substituents;

R⁹ is independently selected from:

- 20 (1) unsubstituted or substituted C₁-C₁₀ alkyl,
- (2) unsubstituted or substituted heterocycle, and
- (3) unsubstituted or substituted aryl;

W is selected from:

- 25 (1) aryl, and
- (2) heterocycle;

m is 0, 1, or 2;

n is independently 0, 1, 2, 3, 4, 5, or 6;


30 p is 0, 1, 2, 3, or 4;

q is independently 0, 1, or 2;

t is independently 0, 1, 2, 3, 4, 5, or 6;

35 or a pharmaceutically acceptable salt, hydrate or stereoisomer thereof.

2. The compound, according to Claim 1, as illustrated by Formula I:



wherein

5 R^{1a} is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C_1 - C_6 alkyl, and
- (3) OR^8 ;

10 R^1 is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C_1 - C_{10} alkyl,
- (3) halo,
- (4) CF_3 ,
- 15 (5) $-(CH_2)_t R^9 C(O)R^8$,
- (6) $-C(O)R^9$,
- (7) $-(CH_2)_t OR^8$,
- (8) $-(CH_2)_t C(O)NR^7 R^8$,
- (9) $-C(O)OR^8$,
- 20 (10) $-(CH_2)_t NR^7 R^8$, and
- (11) $-(CH_2)_t S(O)_q (CH_2)_t NR^7 R^8$;

R^2 is independently selected from:

- (1) H,
- 25 (2) unsubstituted or substituted C_1 - C_{10} alkyl,

- 5 (3) unsubstituted or substituted aryl,
(4) unsubstituted or substituted heterocycle,
(5) unsubstituted or substituted C₃-C₁₀ cycloalkyl,
(6) unsubstituted or substituted C₂-C₆ alkenyl,
(7) unsubstituted or substituted C₂-C₆ alkynyl,
(8) CN,
(9) halo,
(10) N(R⁸)₂, and
(11) OR⁸;

10

R³ is independently selected from:

- (1) H,
(2) unsubstituted or substituted C₁-C₁₀ alkyl, and
(3) unsubstituted or substituted aralkyl;

15

R⁷ is selected from:

- (1) H,
(2) unsubstituted or substituted C₁-C₁₀ alkyl, and
(3) unsubstituted or substituted aralkyl;

20

R⁸ is independently selected from:

- (1) H,
(2) unsubstituted or substituted C₁-C₁₀ alkyl, and
(3) unsubstituted or substituted aryl;

25

R⁷ and R⁸, when attached to the same nitrogen atom may be joined to form a 5-7 membered heterocycle containing, in addition to the nitrogen, one or two more heteroatoms selected from N, O, or S, said heterocycle being optionally substituted with one to three R² substituents;

30

R⁹ is independently selected from

- (1) unsubstituted or substituted aryl, and
(2) unsubstituted or substituted heterocycle;

W is selected from:

- (1) aryl, and
- (2) heteroaryl, selected from pyridyl, pyrimidinyl, isoxazolyl, or pyrazinyl;

5 m is 0, 1, or 2;

n is independently 0, 1, 2, 3, 4, 5, or 6;

p is 0, 1, 2, 3, or 4;


q is independently 0, 1, or 2;

t is independently 0, 1, 2, 3, 4, 5, or 6;

10

or a pharmaceutically acceptable salt, hydrate or stereoisomer thereof.

3. The compound, according to Claim 1, as illustrated by Formula II:



15

wherein

R¹ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- 20 (3) halo,
- (4) unsubstituted or substituted aryl,
- (5) unsubstituted or substituted heterocycle,
- (6) CF₃,
- (7) -(CH₂)_t R⁹C(O)R⁸,
- 25 (8) -C(O)R⁹, and
- (9) -(CH₂)_tOR⁸;

R² is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl,
- (3) unsubstituted or substituted aryl,
- 5 (4) unsubstituted or substituted heterocycle,
- (5) halo,
- (6) OR⁸,
- (7) N(R⁸)₂, and
- (8) CN;

10

R³ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
- 15 (3) unsubstituted or substituted aralkyl;

15

R⁸ is independently selected from:

- (1) H,
- (2) unsubstituted or substituted C₁-C₁₀ alkyl, and
- 20 (3) unsubstituted or substituted aryl;

20

R⁹ is independently selected from

- (1) unsubstituted or substituted aryl, and
- (2) unsubstituted or substituted heterocycle;

25 m is 0, 1, or 2;

n is 0, 1, 2, 3, 4, 5, or 6;

p is 0, 1, 2, 3, or 4;

t is independently 0, 1, 2, 3, 4, 5, or 6;

30 or a pharmaceutically acceptable salt, hydrate or stereoisomer thereof.

4. A compound selected from:

4-(2-amino-5-bromo-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine;


4-(2-amino-1,3-thiazol-4-yl)-N-(3,5dimethylphenyl)pyrimidin-2-amine;

4-(2-amino-5-phenyl-1,3-thiazol-4-yl)-N-(3,5-dimethylphenyl)pyrimidin-2-amine;
 2-amino-4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-
 carbonitrile;
 4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile;


5

or a pharmaceutically acceptable salt or hydrate thereof.

5. The compound according to Claim 4 which is selected from



10 2-amino-4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-
 carbonitrile;



4-{2-[(3,5-dimethylphenyl)amino]pyrimidin-4-yl}-1,3-thiazole-5-carbonitrile;

15 or a pharmaceutically acceptable salt or hydrate thereof.

6. A pharmaceutical composition which is comprised of a
 compound in accordance with Claim 1 and a pharmaceutically acceptable carrier.

7. A method of treating or preventing cancer in a mammal in need of such treatment which is comprised of administering to said mammal a therapeutically effective amount of a compound of Claim 1.

5 8. A method of treating cancer or preventing cancer in accordance with Claim 7 wherein the cancer is selected from cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung.

9. A method of treating or preventing cancer in accordance with
10 Claim 7 wherein the cancer is selected from histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.

10. A method of treating or preventing a disease in which
15 angiogenesis is implicated, which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.

11. A method in accordance with Claim 10 wherein the disease is an ocular disease.

20

12. A method of treating or preventing retinal vascularization which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of compound of Claim 1.

13. A method of treating or preventing diabetic retinopathy which
25 is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of compound of Claim 1.

14. A method of treating or preventing age-related macular degeneration which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.

5 15. A method of treating or preventing inflammatory diseases which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.

10 16. A method according to Claim 15 wherein the inflammatory disease is selected from rheumatoid arthritis, psoriasis, contact dermatitis and delayed hypersensitivity reactions.

15 17. A method of treating or preventing a tyrosine kinase-dependent disease or condition which comprises administering a therapeutically effective amount of a compound of Claim 1.

18. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier.

20 19. A process for making a pharmaceutical composition which comprises combining a compound of Claim 1 with a pharmaceutically acceptable carrier.

25 20. A method of treating or preventing bone associated pathologies selected from osteosarcoma, osteoarthritis, and rickets which comprises administering a therapeutically effective amount of a compound of Claim 1.

21. The composition of Claim 6 further comprising a second compound selected from:

30 1) an estrogen receptor modulator,

- 2) an androgen receptor modulator,
- 3) retinoid receptor modulator,
- 4) a cytotoxic agent,
- 5) an antiproliferative agent,
- 5 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor, and
- 10) another angiogenesis inhibitor.

10

22. The composition of Claim 21, wherein the second compound is another angiogenesis inhibitor selected from the group consisting of a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP
15 inhibitor, an integrin blocker, interferon- α , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, and an antibody to VEGF.

20

23. The composition of Claim 21, wherein the second compound is an estrogen receptor modulator selected from tamoxifen and raloxifene.

25

24. A method of treating cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with
25 radiation therapy.

30

25. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a compound selected from:

- 1) an estrogen receptor modulator,

- 5
- 2) an androgen receptor modulator,
 - 3) retinoid receptor modulator,
 - 4) a cytotoxic agent,
 - 5) an antiproliferative agent,
 - 6) a prenyl-protein transferase inhibitor,
 - 7) an HMG-CoA reductase inhibitor,
 - 8) an HIV protease inhibitor,
 - 9) a reverse transcriptase inhibitor, and
 - 10) another angiogenesis inhibitor.

10

26. A method of treating cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy and a compound selected from:

- 15
- 1) an estrogen receptor modulator,
 - 2) an androgen receptor modulator,
 - 3) retinoid receptor modulator,
 - 4) a cytotoxic agent,
 - 5) an antiproliferative agent,
 - 6) a prenyl-protein transferase inhibitor,
 - 20 7) an HMG-CoA reductase inhibitor,
 - 8) an HIV protease inhibitor,
 - 9) a reverse transcriptase inhibitor, and
 - 10) another angiogenesis inhibitor.

25

27. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 and paclitaxel or trastuzumab.

28. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 and a GPIIb/IIIa antagonist.

5 29. The method of Claim 28 wherein the GPIIb/IIIa antagonist is tirofiban.

30. A method of reducing or preventing tissue damage following a cerebral ischemic event which comprises administering a therapeutically effective
10 amount of a compound of Claim 1.

31. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a COX-2 inhibitor.

15 32. A method of treating or preventing preeclampsia which comprises administering a therapeutically effective amount of a compound of Claim 1.

33. A method of inhibiting at least two tyrosine kinase receptors
20 which comprises administering a therapeutically effective amount of a compound according to Claim 1.

34. The method according to Claim 33 wherein the tyrosine kinase receptors are selected from KDR, EGFR and SRC.

25 35. A method of treating or preventing tissue damage due to bacterial meningitis which comprises administering a therapeutically effective amount of a compound of Claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/23882

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 239/28, 239/48; A61K 31/506; A61P 35/00, 19/02

US CL : 544/321, 324; 514/269, 272, 275

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/321, 324; 514/269, 272, 275

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE, EAST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,958,934 A (BERGER et al) 28 September 1999 (28.09.1999), see entire document, especially see Formula I, column 2, R3 definition.	1-9, 18-19, 25-35

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>		<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	--	---

Date of the actual completion of the international search	Date of mailing of the international search report
11 September 2002 (11.09.2002)	19 SEP 2002
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703)305-3230	Authorized officer <i>Venkataraman Balasubramanian</i> Telephone No. (703)308-1235

Form PCT/ISA/210 (second sheet) (July 1998)

THIS PAGE BLANK (USPTO)